

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES


Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 035 209 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
13.09.2000 Bulletin 2000/37

(51) Int Cl.7: C12N 15/86, C12N 7/01,
C12N 5/10, A61K 39/00,
A61K 39/145, A61K 48/00

(21) Application number: 99104519.6

(22) Date of filing: 06.03.1999

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE
Designated Extension States:
AL LT LV MK RO SI

• Flick, Ramon
13793 Västerhaninge (SE)
• Menke, Anette
35037 Marburg (DE)
• Azzey, Mayra
35394 Giessen (DE)

(71) Applicant: ARTEMIS Pharmaceuticals GmbH
51063 Köln (DE)

(74) Representative:
Helbing, Jörg, Dr. Dipl.-Chem. et al
Patentanwälte
von Kreisler-Selting-Werner,
Postfach 10 22 41
50462 Köln (DE)

(72) Inventors:
• Hobom, Gerd
35392 Giessen (DE)

(54) Stable recombinant influenza viruses free of helper viruses

(57) The invention relates to a recombinant influenza virus for high-yield expression of incorporated foreign gene(s), which is genetically stable in the absence of any helper virus; wherein

- (a) at least one of the regular viral RNA segments has been exchanged for a vRNA encoding a foreign gene; and/or
- (b) at least one of the regular viral RNA segments is an ambisense RNA molecule, containing one of

the standard viral genes in sense orientation and a foreign, recombinant gene in antisense orientation, or *vice versa*, in overall convergent arrangement.

The present invention further provides a method for the production of said recombinant influenza virus, a method for constructing specific ribozyme-sensitive influenza carrier strains; pharmaceutical compositions comprising said recombinant influenza virus; and the use of said recombinant influenza virus for preparing medicaments for vaccination purposes.

EP 1 035 209 A1

Description

[0001] The invention relates to a recombinant influenza virus for high-yield expression of incorporated foreign gene(s), which is genetically stable in the absence of any helper virus; a method for the production of said recombinant influenza virus, a method for constructing specific ribozyme-sensitive influenza carrier strains; pharmaceutical compositions comprising said recombinant influenza viruses; and the use of said recombinant influenza virus for preparing medicaments for vaccination purposes.

[0002] Redesigning influenza virus into a vector system for expression of foreign genes similar to what has been achieved in several other thoroughly studied viruses such as adenovirus, retrovirus, Semliki Forest virus or Rabies virus has the advantage of an industrially well established mode of cheap propagation for influenza in fertilized chicken eggs leading to rather high titers (above 10^{10} /ml). On the other hand none of its genes may be deleted from the influenza genome according to our present knowledge, and give room for large-size foreign insertions. Only small fragments of foreign polypeptide chains such as B cell epitopes (10 to 15 amino acids) may be inserted into selected positions within two of the viral proteins, i.e. in exchange for one of the variable antigenic regions located at the surface of hemagglutinin (Muster et al., Mucosal model of immunization against human immunodeficiency virus type 1 with a chimeric influenza virus. *J. Virol.* 69 (11), 6678-6686 (1995)), or into the stalk sequence of viral neuraminidase (Garcia-Sastre and Palese, The cytoplasmic tail of the neuraminidase protein of influenza A virus does not play an important role in the packaging of this protein into viral envelopes. *Virus Res.* 37, 37-47 (1995)), and be stably maintained as functional fusion proteins. Constructs of this kind turned out to be useful for experimental vaccination in a few cases studied, but only rather few clearly defined epitope sequences (of ten to twelve amino acids each) are known today, and some of them might also be misfolded within such restricted fusion protein positions, or in other cases interfere with the correct tertiary structure and function of their host polypeptide chains.

[0003] Incorporation of a full-size foreign protein into influenza via reverse genetics, encoded by an independent ninth vRNA molecule in addition to its regular set of eight standard vRNA segments is without special provisions only transiently possible (Luytjes et al., Amplification, expression, and packaging of a foreign gene by influenza virus. *Cell* 59, 1107-1113 (1989); Enami et al., An influenza virus containing nine different RNA segments. *Virology* 185, 291-298 (1991)). In the absence of a continuous selective pressure any additional recombinant vRNA segment cannot be stably maintained as long as a wildtype promoter sequence is used on that ninth vRNA segment, and it will inadvertently be lost after few steps of propagation. Using a different system of influenza reverse genetics developed in our laboratory (Zobel et al., RNA polymerase I catalysed transcription of insert viral cDNA. *Nucleic Acids Res.* 21, 3607-3614 (1993); Neumann et al., RNA polymerase I-mediated expression of influenza viral RNA molecules. *Virology* 202, 477-479 (1994)), which was built around *in vivo* synthesis of recombinant vRNA molecules by cellular RNA polymerase I transcription of the respective template cDNA constructs, promoter-up mutations have been designed by nucleotide substitutions (Neumann and Hobom, Mutational analysis of influenza virus promoter elements *in vivo*. *J. Gen. Virol.* 76, 1709-1717 (1995)). When these are attached to a recombinant ninth vRNA segment its increased transcription and amplification rate will not only compensate for the losses suffered spontaneously, but even cause accumulation of the foreign vRNA segment during simple viral passaging, in the absence of any selection. However, due to its over-replication relative to all of the regular influenza vRNA segments (which of course are connected to wild-type promoter sequences) after catching up with the others the foreign segment will become over-abundant. This increasingly will result in viral particles that have incorporated several copies of recombinant vRNA, but no longer have a full set of all eight standard segments incorporated among an average of about 15 vRNA molecules present within a virion. Such particles are defective and will not result in plaque formation, hence after an initial increase of recombinant viral particles during the first steps of propagation a dramatic decrease is observed, usually at the third or fourth step of viral passaging, depending on the size of the recombinant vRNA and the level of the promoter-up mutation attached. A balanced situation with regard to the insert length and the level of promoter activity can be achieved, and has been propagated in a particular case over 11 passages, with essentially stable levels of recombinant viruses among a majority of helper viruses (around 80%) during these steps. If a full-level promoter-up mutation is used (1104 or the new variant 1920, see below) a balanced-level propagation is reached in conjunction with a recombinant vRNA size of 4000 nucleotides (unpublished):

[0004] In all of these preparations, both in transiently achieved increased yields (up to 40% of recombinants after three or four steps of viral passage), and in a balanced propagation of recombinant influenza viruses (10 - 20%) the respective viral progeny inadvertently constitute mixtures with a majority of non-recombinant helper viruses. These result both from a statistical mode of packaging vRNA molecules into a virion (the ninth segment may not be co-packaged), and from the fraction of cells solely infected by helper virus. The problems of fractional yields and of instability during viral propagation of recombinant influenza are the problems to be solved with the present invention.

[0005] Starting out from two observations in this laboratory which are discussed below and which concern two hitherto unsuspected properties of influenza viral RNA polymerase in its interaction with terminally adapted influenza-specific RNA molecules, a new technique for the construction of stable recombinant influenza viruses was found.

[0006] As previously described in WO 96/10641 plasmid constructs are designed to generate influenza vRNA-like molecules *in vivo* by cellular RNA polymerase I transcription following plasmid DNA transfection into tissue culture cells, and to this end contain flanking rDNA promoter and terminator elements, externally located relative to any cDNA constructs in between. The resulting recombinant vRNA molecules are designed to contain 5' and 3' recognition sequences for influenza viral RNA polymerase, which however carry up to five nucleotide substitutions (in promoter-up mutant pHL1920) resulting in a substantial increase of expression over wildtype influenza promoter levels. While recombinant pseudoviral RNA is initially transcribed by RNA polymerase I, further amplification and mRNA transcription depends on the presence of viral RNA polymerase and viral nucleoprotein in the cell, which generally are provided by infection of a helper virus. As a consequence the progeny viral yield will constitute a mixture of recombinant viruses together with a majority of wild-type helper viruses.

[0007] In the new technique the recombinant vRNA-like molecules as transcribed by RNA polymerase I are constructed as ambisense RNA double segments, with one reading frame (an influenza gene) in sense and a second one (a foreign gene) in anti-sense orientation, or vice versa. In such constructs both reading frames will be expressed via the cap-snatching mode, even if at different levels. Again, infection by helper virus is required to provide the necessary viral early and late proteins for genetic expression and virion packaging. However, the particular helper virus used in the new method is a recombinant virus carrying 2x2 specifically designed ribozyme targets inserted into the flanking non-coding sequences of one of its eight vRNA segments. The viral segment chosen to become ribozyme-sensitive is always identical to the one used in constructing the recombinant ambisense RNA molecule, as the viral carrier gene in covalent linkage with an additional foreign gene.

[0008] Recombinant influenza viruses produced in this way through RNA polymerase I transcription of an ambisense viral RNA molecule followed by infection with that specifically designed type of ribozyme-sensitive helper virus will carry one of the influenza genes twice, once within that ribozyme-sensitive helper vRNA segment, and a second time within the ribozyme-resistant ambisense segment. Recombinant viruses of this type are again obtained initially only as a mixture together with a majority of non-recombinant helper viruses. A progeny viral passage through tissue culture cells (293T) which before have been transiently transfected with plasmid constructs expressing the respective double-headed ribozyme will (in one step) inactivate the ribozyme-sensitive segment by a factor of up to 100. One or two rounds of such ribozyme treatment *in vivo* will at the same time (a) purify the recombinant virus from its non-recombinant helper contaminants, and (b) delete the sensitive vRNA helper segment from within the initial (additive) recombinant virus.

[0009] As a result recombinant influenza viruses are isolated along this several-step procedure, which are free of contaminating helper viruses and carry seven regular and one ambisense vRNA segments, all in a balanced replication mode. Their recombinant nature is stably maintained because of a covalent junction between one of the viral genes and the full-size foreign gene inserted, a situation achieved here for the first time, via constructing an influenza ambisense RNA segment. The whole procedure is independent of any (selectable) phenotype, and can be applied to either of the eight influenza vRNA segments. After establishing a first ambisense vRNA segment carrying a single foreign gene it can also be repeated all over for inserting a second foreign gene within another ambisense RNA segment of the same constitution in principle.

[0010] Stable recombinant viruses of the type described can be used for cheap propagation in fertilized eggs, either for production of those recombinant viruses themselves or for production of foreign proteins or glycoproteins encoded by them, and hence find application in (glyco)protein production or in providing vector systems for somatic gene therapy or in being used as vaccination agents.

[0011] Thus, the present invention provides

(1) a recombinant influenza virus for high-yield expression of incorporated foreign gene(s), which is genetically stable in the absence of any helper virus, wherein

(a) at least one of the regular viral RNA segments has been exchanged for a vRNA encoding a foreign gene; and/or

(b) at least one of the regular viral RNA segments is an ambisense RNA molecule, containing one of the standard viral genes in sense orientation and a foreign, recombinant gene in anti-sense orientation, or *vice versa*, in overall convergent arrangement;

(2) a method for the production of recombinant influenza viruses as defined in (1) above comprising

(a) RNA polymerase I synthesis of recombinant vRNAs *in vivo*, in antisense, sense or ambisense design,
 (b) followed by infection with an influenza carrier strain constructed to include flanking ribozyme target sequences in at least one of its viral RNA segments, and
 (c) thereafter selective vRNA inactivation through ribozyme cleavage;

(3) a method of constructing influenza carrier strains carrying one or more ribozyme target sites (of type one) in vRNA flanking positions comprising

(a) RNA polymerase I synthesis of recombinant vRNAs *in vivo*, carrying two different 3' promoter sequences in tandem, which are separated by a second type of ribozyme target sequence, and which carry the said internal ribozyme target sites of type one;

(b) followed by infection of an influenza wildtype strain;

(c) thereafter amplification through simple steps of viral propagation; and

(d) finally isolation through removal of their external 3' promoter sequence by ribozyme cleavage through infection of cells expressing ribozyme type 2, followed by plaque purification;

said method being suitable for the construction of an influenza carrier strain required for step (b) of (2) above;

(4) a ribozyme-sensitive influenza carrier (helper) strain obtainable by the method of (3) above;

(5) a pharmaceutical composition comprising a recombinant virus as defined in (1) above;

(6) use of a recombinant virus as defined in (1) above for preparing a medicament for vaccination purposes;

(7) use of a recombinant virus as defined in (1) above as vector systems in somatic gene therapy, for transfer and expression of foreign genes into cells (abortively) infected by such viruses, either in *ex vivo* or *in vivo* application schemes; and

(8) a method for the production of proteins or glycoproteins which comprises utilizing a recombinant influenza virus as defined in (1) above as expression vector.

[0012] Moreover, the invention provides a method for preventing/treating influenza utilizing a recombinant virus as defined in (1) above, i.e., a vaccination method utilizing said recombinant virus.

[0013] According to the present invention "influenza virus" embraces influenza A virus, influenza B virus and influenza C virus. A preferred recombinant virus of the invention is where one or both of the standard glycoproteins hemagglutinin and neuraminidase have been exchanged into foreign glycoprotein(s) or into fusion glycoproteins consisting of an anchor segment derived from hemagglutinin and an ectodomain obtained from the foreign source, viral or cellular, or in which such recombinant glycoprotein has been inserted as a third molecular species in addition to the remaining standard components.

[0014] Further preferred recombinant viruses of the invention are where the terminal viral RNA sequences, which are active as the promoter signal, have been varied by nucleotide substitutions in up to five positions, resulting in improved transcription rates of both the vRNA promoter as well as the cRNA promoter as present in the complementary sequence.

[0015] Further preferred embodiments of the invention are set forth herein below.

A. Construction of influenza helper virus strains carrying ribozyme target sequences in flanking positions within either of the vRNA segments

[0016] A.1. Influenza RNA polymerase will initiate transcription and replication from promoter structures located at internal positions in an RNA molecule, not only from the natural position at both ends of a vRNA molecule:

[0017] This is true in particular for promoter-up variants in RNA-internal location due to their enhanced binding affinity for viral polymerase. Not only 3' end extensions are tolerated in RNA-internal promoter recognition (Fig. 1), but also 5' extensions as well as extensions at both ends of the RNA template molecule, containing noncomplementary as well as complementary sequence, i.e. potentially present as a double-stranded extension. Finally, also an extension by way of duplication of the promoter sequence (active or inactivated) leads to mRNA transcription and CAT expression, initiated from the active pair of 5' and 3' promoter halves, irrespective if in external or in internal or even in an oblique localization (active 5' promoter sequence in external, active 3' promoter sequence in internal position, or vice versa). RACE-determination of the resulting 5' and 3' ends of viral mRNA and cRNA, i.e. the products of transcription and replication reactions for several of the extended template vRNA constructs proves an exact recognition and sequence-specific initiation at a position equivalent to regular 3' position 1: all of the various template extensions are lost in every product RNA molecule, most likely after only one round of replication.

[0018] A.2. Bicistronic (tandem) vRNA molecules carrying an additional 3' specific promoter sequence in a central position between its two genes can be used for an indirect selection method for recombinant influenza viruses:

[0019] The method described is applicable for any foreign gene (e.g. CAT) without a selection potential of its own, if inserted into the distal mRNA position (proximal vRNA position, in anti-sense orientation) behind a carrier gene (e.g. GFP) in the proximal mRNA position, able to serve as a transient selection marker. The carrier gene which is used for selection will get lost spontaneously during further propagation. These constructs are equivalent to a 3' extension of the template vRNA by a full-size gene of 750 nucleotides up to a second 3' promoter sequence, in terminal location.

While in the set of experiments shown in Fig. 2 the external 3' promoter sequence is maintained throughout as the same promoter-up variant (1104), the internal 3' promoter sequence has been varied to include a full-level promoter-up variant (pHL2270, containing promoter mutant 1104), a medium-level promoter variant (pHL2350, promoter mutant 1948), a wildtype promoter construct (pHL2629), and a construct carrying an unrelated central sequence in an otherwise identical design (pHL2300).

[0020] Due to the presence of two 3' promoter sequences in conjunction with a single 5' promoter sequence an alternating interaction between them will constitute one or the other active promoter structure (see Fig. 3). While the external location with an adjacent RNA 3' end may have a structural advantage, this appears to be compensated by the shorter distance in an interaction between the 5' sequence and the central 3' sequence in constituting the internal promoter, such that the competition between the two primarily reflects the various internal 3' promoter allele sequences used, compare Fig. 2B and activity ratios indicated above and below the lanes. Translation of the mRNA-distal gene (CAT) is only observed following an internal initiation at the bicistronic vRNA template, resulting in a spontaneous deletion of the mRNA-proximal (vRNA-distal) gene, GFP, compare right half of Fig. 3. In complementary analyses GFP fluorescence is observed initially for all of the bicistronic constructs, but gets lost on a faster rate from pHL2270 transfected and influenza infected cells (not shown), and will stay unchanged in pHL2300-treated cells. The indirect selection system based on bicistronic (tandem) molecules as designed here and demonstrated for reporter genes GFP and CAT can be used for any other gene without distinct phenotype upon insertion behind an unrelated carrier gene with properties useful in selection. - In employing that technique an initial phase of (repeated) positive selection for infected cells expressing that proximal trait (e.g. by FACS or by magneto-beads) will be followed by a second phase with negative selection, i.e. against that fraction of infected cells still exposing the same property.

[0021] A.3. Isolation of an influenza strain designed to carry 2x2 flanking ribozyme target sequences at the 5' and 3' end of vRNA segment 4 coding for hemagglutinin:

The above scheme for an indirect selection of any foreign recombinant gene behind a proximal carrier gene is further modified by deleting the carrier gene altogether. Instead, both 3' terminal promoter sequences (mutant and wildtype) follow each other at a short distance, separated only by a specific, repetitive ribozyme target sequence, - different from other target sequences to be described further below. The cDNA insert following after the second 3' promoter sequence consists mainly of a regular hemagglutinin (H7) coding sequence, however both the 5' and 3' vRNA terminal regions of the insert carry that other ribozyme target sequence (different from the first target sequence mentioned above) inserted in either location in tandem duplication (pHL2969, see Fig. 14).

[0022] Due to a superior replication supported by the promoter-up variant located in 3' external position the recombinant HA segment attached to that promoter sequence will become enriched during the first steps of viral propagation, while the originally dominating HA segment of the helper virus which is under control of a wildtype promoter sequence is consecutively reduced and finally is no longer detectable among viral progeny. This result is documented by RT-PCR analysis of consecutive viral populations as obtained in that stepwise propagation procedure, see Fig. 4.

[0023] In the next step the viral lysate is twice passaged via infection of cell culture cells (293T) that before have been DNA-transfected by plasmid pAM403 (Fig. 15). This construct has been specifically designed to express a hammerhead ribozyme with flanking sequences complementary to the repetitive GUC-containing target sequence, as present twice in between the external and internal 3' promoter signals in the recombinant HA vRNA segment, see Fig. 5. In this way the extra promoter sequence is cut off from the finally resulting recombinant HA vRNA. Its promoter-up activity was useful in achieving an initial increase in the concentration of recombinant HA vRNA over wildtype HA vRNA, and in finally excluding the latter from further propagation. However, for the same reason that high activity of the promoter variant will cause instability in the resulting viral progeny, and an effective 'substitution' at this time through ribozyme cleavage by the internally located promoter signal, wild-type or slightly improved, will restore stability to the progeny viruses, with all of their eight vRNA segments now brought back in balance to each other. Due to the ribozyme cleavage site at 26 nucleotides 3' of the wild-type promoter sequence (see Fig. 5), in the initial stage that promoter signal is situated in a vRNA-internal location, extended by a 3' terminal sequence of 26 nucleotides. According to the data presented in Fig. 1 this should cause a transient slight reduction in activity, resulting however in one step in regular viral mRNAs and cRNAs, with any initially remaining extra sequence being lost from the finally resulting recombinant HA vRNA.

[0024] Progeny viruses still carrying an external promoter-up sequence (before ribozyme treatment or due to incomplete reaction) will not cause any plaque, due to over-replication of one vRNA segment relative to all others which results in a high load of defective particles. However, progeny viruses which have lost that external promoter element due to ribozyme cleavage will yield regular plaques due to a balanced mode of replication for all eight wild-type or recombinant vRNA segments. Hence plaque purification is used for isolating a pure influenza viral strain carrying 2x2 ribozyme targets in its recombinant HA vRNA segment, with its termini reduced to the wild-type promoter sequence. The nature of the viral strain isolated has been confirmed in this regard by RT-PCR analysis, see Fig. 4.

[0025] The above isolation procedure resulting in influenza viral strains carrying 2x2 flanking ribozyme target sequences has been performed twice for the HA coding segment (segment 4) to obtain two different isolates with regard

to the orientation of the ribozyme target sequences. In one of the isolates (vHM41, see SEQ. ID NO: 3) the tandem target sites have been inserted into the HA vRNA non-translated sequence both in 5' and 3' location, while according to the second design that 5' tandem target sequence has been included in an inverted orientation, such that it is now present in the cRNA 3' sequence instead (vHM42).

[0026] In another experiment the same procedure was used to isolate an influenza strain carrying 2x2 tandem target sites within the 5' and 3' flanking positions of segment 8 vRNA, i.e. coding for genes NS1 and NS2 (vHM81; see SEQ. ID NO:4). And in principle the same could be done for any other influenza segment, in particular since only the reading frame cDNA sequence has been exchanged from HA to NS, with all of the flanking elements directly responsible for that procedure remaining in place, unchanged.

[0027] A.4. Ribozyme cleavage and vRNA segment exchange using ribozyme-sensitive influenza strains as helper viruses:

In an initial model experiment a range of ribozyme type and target site locations was probed in designing a series of CAT reporter gene vRNA constructs (analysed in the presence of a surplus of wildtype helper virus) in 293T cells. While all of the ribozyme constructs adhered to the hammerhead model, with 10 to 12 nucleotides of complementary

sequence flanking on either side of the GUC target site, these ribozyme constructs varied from monomer to dimer to trimer repetitions. Ribozyme containing mRNAs were synthesized *in vivo* via the basic vector plasmid pSV2-*neo*, i.e. using the efficient p_{SVe} RNA polymerase II promoter element for expression, and the SV40 origin signal for plasmid amplification, in a cell line (293T or cos-1) with an incorporated SV40 T antigen gene. In addition the pSV2-*neo* mRNA includes the small, 63 nucleotide intron sequence of the SV40 early mRNA which is supposed to be spliced very slowly, thereby extending the pre-mRNA half-life in the nucleus. Each of the pSV2-*neo*-ribozyme plasmid constructs was transfected into 293T cells. Thereafter, recombinant viruses containing dimer target sites either near one end of the molecule only, or near both ends have been used for infection of the transfected cells. Relative activities of ribozyme constructs versus vRNA target sites have been measured via inactivation of CAT acetyl transfer rates in the cell lysates obtained at 8 h post infection (Fig. 6). The highest activities were observed for dimer ribozymes acting on vRNA molecules carrying 2x2 target sequences on both ends of the molecule, either in vRNA 5' and 3' location, or in vRNA 3' and cRNA 3' location, i.e. with an inversion of the target site sequence at the vRNA 5' end. Consequently, the two constructs carrying tandem ribozyme double targets within both of their non-translated vRNA flanking sequences have been used in the design of ribozyme-sensitive influenza virus strains as described above, with both variants isolated for segment 4 (HA), and one of them for segment 8 (NS). In complementary correspondence the hammerhead ribozyme plasmid used has also been constructed as a double-headed structure with flanking sequences as shown in Fig. 7 (pAM424; for its detailed structure, see Fig. 16).

[0028] The three target site-containing influenza strains isolated as described above have been analysed for their sensitivity against ribozyme cleavage by infection of 293T cells, which had been DNA-transfected 18 h earlier by ribozyme-producing plasmid pAM424, at DNA-transfection rates between 60 and 70 % (as observed in parallel transfections using GFP-expressing plasmid pAM505). Inactivation rates in these one-step control experiments were between 90% and 99% for all three ribozyme-sensitive strains, in their extent mainly depending upon the actual DNA-transfection rates achieved in individual experiments.

In the next step both isolates of HA-coding ribozyme-sensitive viruses, vHM41 and vHM42, have been used in marker-rescue experiments. Here, 293T cells have been first DNA transfected by HA-variant cDNA construct pHL2507 (see Fig. 17), followed after 18 h by vHM41 or vHM42 virus infections at moi 1. The resulting viral supernatant containing e.g. a mixture of ribozyme-sensitive vHM41 carrier virus and pHL2507/vHM41 recombinant virus is propagated in an intermediate step on MDCK cells, which also results in an increase in the fraction of recombinant viruses. Thereafter the resulting virus-containing supernatant is passaged through 293T (or cos-1) cells, which in advance have been transiently transfected by ribozyme-producing pAM424. As may be concluded from the above experiment (Fig. 6) and shown in Fig. 8 vHM41- or vHM42-derived ribozyme-sensitive HA vRNA segments are expected to be inactivated by pAM424-produced ribozyme down to a remaining level of about 1% to 10% (mainly present within cells that are infected, but not DNA-transfected).

[0029] Instead, the substitute HA vRNA which originated from pHL2507 plasmid DNA (ribozyme-resistant) becomes exclusively incorporated into progeny virions. For further purification and viral propagation these have been passaged a second time through 293T cells, again in advance DNA-transfected by pAM424, and after another amplification step on MDCK cells the resulting viral preparations have been used for RT-PCR analysis. The resulting viral populations in these marker rescue experiments only contain HA-vRNA molecules derived from pHL2507, which in their PCR analyses are of intermediate size relative to wildtype HA-vRNA, and vHM41- or vHM42-derived HA-vRNAs, respectively.

[0030] Consequently, a set of ribozyme-sensitive influenza strains with targets inserted individually into every vRNA molecule may be used for such one-step marker rescue experiments in general, i.e. for vRNA segment exchange reactions performed in a directional way for any of the eight influenza vRNA segments, without requirement for a selectable change in phenotype (genetic marker).

B. Expression of two gene products from ambisense bicistronic influenza vRNA

[0031] B.1. The influenza cRNA promoter is active in antisense viral mRNA transcription according to the cap-snatching mode of initiation:

While the vRNA template of influenza virus is known to be active in viral mRNA as well as cRNA synthesis, the cRNA template has been described so far only to produce vRNA molecules, as a second step in viral replication. The potential activity of the cRNA promoter in initiating also viral mRNA transcription has not been analysed or even suspected so far, since no antisense (vRNA) reading frame can be detected in any of the viral RNA segments. Also, no U_5/U_6 template sequence element is present in any of the viral cRNAs in an adjacent position to its 5' promoter structure as is the case for all viral vRNAs. This element is known to serve as a template sequence for mRNA terminal poly-adenylation, in repetitive interaction. However, when both elements are provided through reconstruction of an artificial influenza cRNA segment: a reading frame in opposite orientation (CAT), and a U_6 template element in 5' adjacent location, CAT expression can indeed be observed, see pHL2583 (see Fig. 18) in Fig. 9. Similar to the vRNA promoter the cRNA promoter activity is improved by (the same) promoter-up mutations, which essentially consist of basepair exchanges according to the 'corkscrew' model. This model apparently also holds for the cRNA promoter structure as analysed in a stepwise manner in Fig. 9. While the cRNA promoter has to be superior over the vRNA promoter in its initiation of replication, since the vRNA/cRNA product ratio was determined to be around 10:1 (Yamanaka et al., *In vivo* analysis of the promoter structure of the influenza virus RNA genome using a transfection system with an engineered RNA. Proc. Natl. Acad. Sci. USA 88, 5369-5373 (1991)), the cRNA promoter is observed to be inferior to the vRNA promoter in its initiation rate of transcription (compare pHL2583 with pHL1844 in Fig. 9), at least for all promoter variants tested so far.

[0032] A RACE analysis for determination of the 5' ends in pHL2583 cRNA promoter transcribed mRNAs proved this initiation to occur according to the cap-snatching mode, in complete equivalence to standard vRNA promoter controlled transcription initiation.

[0033] B.2. Development of ambisense influenza constructs for consecutive expression of two genes (GFP and CAT) from a single viral RNA:

For a bidirectional transcription and translation of influenza RNA segments the two reporter genes GFP and CAT have been arranged in opposite orientation to each other, and the flanking 5' and 3' promoter sequences (adhering to promoter-up variant 1104) had to be reconstructed to include a U_6 poly-adenylation element in either orientation in a 5' promoter adjacent position. This requirement necessarily resulted in a promoter-adjacent 5'- $U_6/3'$ - A_6 complementary structure, both in the vRNA and cRNA terminal sequence (see Fig. 10), which had to be tested for its promoter activity, in either orientation. Therefore, the convergent pair of reporter genes GFP and CAT has been inserted in both orientations, such that CAT transcription is initiated by the vRNA promoter in one construct (pHL2960), and by the cRNA promoter in the other (pHL2989, Fig. 19), and vice versa for GFP expression from both ambisense constructs. In addition, also the CAT gene only has been inserted in either orientation between the 5' and 3' elements of that ambisense promoter, with CAT transcribed by the vRNA promoter in one case (pHL2959), and by the cRNA promoter in the other (pHL2957). The whole set of constructs allows for a direct comparison with corresponding reference constructs carrying a regular vRNA promoter (pHL1844) or cRNA promoter structure (pHL2583), i.e. carrying only the 5'-adjacent U_6 sequence element and no 3'- A_6 counterpart. The two groups of constructs also differ in insert size, since a single inserted gene roughly accounts for 750 nucleotides, the convergent set of two genes for 1500 nucleotides, with the distal half of both mRNAs in this case remaining untranslated, a situation unusual for influenza viral mRNAs.

[0034] As is demonstrated in Fig. 10B for CAT expression of the various ambisense constructs all of them are able to initiate transcription in both orientations, even if at different levels with regard to their vRNA and cRNA promoter-dependent expressions, and also with regard to the insert lengths and convergent arrangements of the GFP/CAT versus CAT-only constructs. Analysis of the GFP expression rates (not shown) yields complementary results, i.e. again vRNA promoter-controlled GFP expression is superior over cRNA promoter expression of GFP. Therefore, individual ambisense clones either show an asymmetric high expression of GFP and low expression of CAT (pHL2960) or vice versa (pHL2989), depending on their orientation of reading frames with regard to the external vRNA and cRNA promoter. Fig. 10C also demonstrates successful propagation of recombinant viruses containing ambisense RNA molecules, which proves survival through amplification, packaging into virions, and expression of both mRNAs in infected MDCK cells (including besides CAT also GFP expression).

[0035] B.3. Construction of a superior promoter-up mutation, pHL1920, to be used for improved rates of cRNA promoter expression in ambisense constructs:

An extended analysis of promoter variants, in particular of complementary double exchanges according to the 'corkscrew' model yielded among others variant pHL1920 (Fig. 20) with CAT activity rates considerably above (125-130% of) the rates observed for standard promoter-up variant '1104' (as present in pHL1844). The '1920' promoter-up variant consists of altogether 5 nucleotide substitutions relative to the wildtype promoter sequence, both in the 5' promoter element (2), and the 3' promoter element (3). The structure of this variant and the whole set of complementary double exchanges is presented in Fig. 11, together with the respective CAT activity measurements, in vRNA promoter con-

structs. vRNA promoter-up variants also show similarly improved expression in (ambisense) cRNA constructs, even if at generally lower levels than in vRNA constructs. cRNA promoter-up expression is observed at levels similar or somewhat (2x-5x) above the *wild-type* vRNA promoter rate, while vRNA promoter-up constructs show CAT expression rates increased up to 20 or 25 times the wild-type vRNA promoter level. In either case expression rates also depend on the size of the insert, with promoter activity rates decreasing with increasing lengths of the influenza RNA molecules to be transcribed.

[0036] B.4. Influenza recombinant viruses containing a foreign gene (CAT) in covalent ambisense linkage with one of the viral genes (HA, NS1/NS2):

The principle solution in designing stable recombinant viruses based on the new properties observed for influenza transcription and replication signals consists in constructing viruses which contain a foreign gene in covalent linkage with one of the (indispensable) viral genes, in ambisense bicistronic organization. Preferably the viral gene is connected to the cRNA promoter, while vRNA promoter expression is used for expression of the foreign gene at rates considerably above the viral mRNA synthesis. The promoter-up variant chosen for constructing the ambisense RNA segment intends to bring its cRNA promoter expression (approximately) into balance with all other viral gene expression levels, which are controlled by wild-type vRNA promoters located at the termini of the seven ordinary influenza segments; the respective choice has to take into consideration the overall length of the ambisense segment.

[0037] Isolation of the ambisense recombinant virus employs an RNA polymerase I-transcribed ambisense cDNA construct, which will give rise *in vivo* to ambisense cRNA-type molecules, see Fig. 12. The plasmid DNA transfection mixture used in this step with 293T cells in addition may or may not contain four 'booster' plasmids which under ρ_{CMV} -control produce the four early influenza proteins from non-viral mRNAs: NP, plus PB1, PB2, and PA, i.e. the three subunits of viral polymerase (Pleschka et al., A plasmid-based reverse genetics system for influenza A virus J. Virol. 70, 4188-4192 (1996)), which will increase in a pre-amplification step the copy number of that ambisense viral cRNA segment. At 18 h post transfection the 293T cells are infected by a ribozyme-sensitive influenza strain, e.g. vHM41, which will supply (again) early and also late viral RNAs. The resulting supernatant which contains a mixture of vHM41 carrier virus and vHM41-derived ambisense recombinant virus (nine vRNA segments) is then passaged directly or via an intermediate step of amplification on MDCK cells onto 293T cells that have in advance been DNA-transfected by ribozyme-producing pAM424. Here, the ribozyme-sensitive vRNA segment of vHM41 will be cleaved at its 2x2 target sites by pAM424 specific ribozymes. In recombinant viruses the vRNA gene lost in this way is re-supplemented through its presence within the ambisense segment. The virus-containing supernatant is passaged for amplification and further purification through ribozyme treatment a second time on 293T cells which again have been pretreated by pAM424 DNA transfection.

[0038] Absence of ribozyme-sensitive vRNA, and presence only of ambisense RNA in RT-PCR analysis at this stage allows for further amplification on MDCK cells and a final virus stock preparation on embryonated chicken eggs. CAT assays can be used to analyse for the presence and monitor the activity of this model foreign gene through the various steps of isolation and propagation as well as document technical improvements that might be worked out for one or more of the processive stages.

C. Examples for application of helper-free, stable recombinant influenza viruses

[0039] C.1. Incorporation of reporter gene GFP in NS/GFP or HA/GFP ambisense segments:

Recombinant viruses of this type will allow to follow-up on influenza infection instantly and continuously in individual infected cells, which may also be counted or documented by fluorescence photography or FACS sorting. With improved temperature resistance (Siemering et al., Mutations that suppress the thermosensitivity of green fluorescent protein. Curr. Biol. 6, 1653-1663 (1993)) GFP expression becomes visible at 2 h after infection and shows bright fluorescence after 4 h p.i., it will be possible to follow-up on the spread of viral infection by GFP fluorescence. Stable fluorescence in abortively infected cells e. g. observed in *ex vivo* treatment of dendritic cells similarly supports a follow-up on their reincorporation into animals; other genes may be incorporated by ambisense vRNA into dendritic cells in the same way.

[0040] C.2. Construction of glycoprotein CSFV-E2 carrying influenza, helper-free:

The glycoprotein E2 of CSF virus has been incorporated previously into influenza both as an HA-anchor fusion protein within the viral envelope, and as an additional, unstable ninth vRNA segment into its genome (Zhou et al.; Membrane-anchored incorporation of a foreign protein in recombinant influenza virions. Virology, 246, 83-94 (1998)). Stabilization is now achieved through an ambisense connection with either of the regular viral RNA segments (NS or HA) which also allowed to reach a level of 100% recombinant viruses instead of an hitherto only 20% (Fig. 13), since all carrier viruses are destroyed through ribozyme action, see Figs. 8 and 12. The helper virus containing preparation has already been used successfully as a vaccine against CSFV infection (antibody titers of 1:40000); the increase achieved in recombinant viruses allows a further improvement in that regard. Also (cost-effective) propagation in fertilized chicken eggs has become possible due to its stable incorporation of the foreign gene as a covalent ambisense construct.

[0041] C.3. Construction of hepatitis C glycoprotein-carrying recombinant influenza viruses as a candidate vaccine:

Hepatitis C virus is a close relative of CSF virus (hog cholera virus), and in particular its set of two glycoproteins, small-size E1 and larger-size E2, is closely related in structural detail and presumably also in function to the corresponding CSFV proteins. An incorporation of HCV-E2/HA fusion proteins into influenza viral envelopes has been achieved in analogy to the CSFV-E2/HA incorporation. In addition, incorporation of an anchor-fusion glycoprotein HCV-E1/HA or both together (in NS and HA ambisense junctions) allows further variations in constructing an influenza-based vaccine for hepatitis C. In analogy to CSFV-E2, neutralizing antibodies are expected to be directed against particular epitopes of HCV-E2, presented in essentially native conformation at the influenza viral envelope.

[0042] C.4. Stable incorporation of selected influenza T-cell epitopes in ambisense constructions:

Influenza infection is known to result in both, antibody production against that specific viral strain or indeed its epitopes that are located mainly at the surface of HA, and in an increase of specifically primed cytotoxic T-lymphocytes, stimulated by T-cell epitopes primarily located within the essentially invariable core structure of the NP protein. While the humoral response will result in life-long immunity against that particular strain of influenza or its epitope structures, the T-cell response will be lost or severely reduced some time afterwards, such that its specificity against influenza in general will fall below protective levels. One way in trying to increase that level of cellular immunity is to enhance the response or recruitment of influenza-specific CTL cells by increasing the level of T-cell epitopes in the infected cell and hence its presentation on the surface by MHC-I receptors. This is achieved by combining in an ambisense construct the HA gene and a series of repeated T-cell epitope sequences as present in the influenza NP gene, in a model design specific for the BALB/C mouse MHC-I allele. Here, the promoter-up expression rate is realized (in vRNA promoter-controlled initiation) for expression of the repetitive epitope polypeptide chain. Alternatively or in addition a controlled secretion of an interleukin can be achieved from recombinant influenza-infected cells, upon ambisense incorporation of the respective gene preferably into the NS segment. The interleukin to be chosen for this purpose (IL-12 or other) is selected to enhance the longevity of influenza-specific CTL cells or its conversion into corresponding memory cells. In this way an ambisense vaccine strain against influenza itself is achieved with expected protective capacity against influenza in general.

[0043] C.5. Exchange of influenza glycoproteins against foreign viral glycoproteins (VSV-G):

The 'marker rescue' experiment described above (section A.4.), i.e. an exchange of one HA gene (ribozyme-sensitive) for another (ribozyme-resistant) can also be used for an exchange of HA for an entirely different glycoprotein, such as the vesicular stomatitis virus G protein, as long as it is attached to the HA anchor segment. Effective incorporation depends always on that C-terminal tail sequence and its interaction with underlying matrix protein M1, and therefore, all constructions consist of fusion proteins in direct analogy to CSFV-E2. VSV-G with or without a foreign anchor sequence has been shown in several other viruses to be able to substitute for the original glycoprotein and to result in infectious viruses with VSV-G specific host-ranges (e.g. in retroviruses, rabies virus, measles virus). G protein in VSV itself as well as on the surface of foreign viruses is the only glycoprotein required for all of the consecutive steps in infection. Insertion of VSV-G instead of HA in recombinant influenza viruses leaves the second glycoprotein, neuraminidase, without any function, which then will get lost spontaneously from the recombinant viruses. This will further increase the capacity for an addition of foreign genes, beyond the gain resulting from an exchange of the larger HA for the smaller VSV-G, which might be used for an addition of ambisense constructs.

[0044] The invention is further illustrated in the accompanying figures.

Description of the figures

[0045]

Fig.1. 3' nucleotide extensions of influenza vRNA template molecules:

(A) Murine B82 cells have been transfected by plasmid cDNA constructs designed to be transcribed into influenza vRNA molecules by RNA polymerase I *in vivo*, followed after 20 h by standard FPV^{Bratislava} helper virus infection, at an moi of 1 to 3. In addition to reference plasmid pHL2024 (no extension), related cDNA constructs carrying extensions of 1 to 50 bp, and hence extended by 1 to 50 nucleotides at the resulting vRNA 3' ends were used in parallel transfections; template extensions are marked at the top of the figure. Cell lysates prepared at 8 h post helper virus infections were used for CAT reactions using in one round 50 µl of cell lysate each, and in further analyses 5 µl and 0.5 µl of lysate (not shown). Relative yields were determined in comparison to reference plasmid pHL2024, as indicated below the figure, with calculations restricted to those CAT assays showing less than 40% of substrate consumption, in three or more independent experiments.

(B) Viral passage of B82 supernatants containing recombinant influenza virus onto MDCK cells, at an moi of 2 to 4, in average. Again at 8 h post infection cell lysates have been prepared and used for CAT assays. Relative yields as indicated below the figure have been determined in comparison to pHL2024 the same way

as in (A), using 0.5 μ l of cell lysate in each case. (The 50 μ l CAT assays as shown here and also in the following figures intend to give an immediate impression of relative activities at always the same level, while the actual measurement data as indicated below the lanes are obtained at various appropriate enzyme concentrations relative to reference pHL2024.)

Fig. 2. Propagation of recombinant influenza viruses with tandem bicistronic vRNA:

(A) General design of expression plasmids for transient bicistronic vRNAs coding for GFP in the mRNA-proximal, and for CAT in the mRNA-distal position. Among the functional elements indicated are the human RNA polymerase I promoter (p_{IH}) and murine rDNA terminator (t_t) sequences, both hatched, and the 5' and 3' vRNA promoter cDNA sequences, open and closed boxes, respectively. For the vRNA-internal 3' promoter signal three variant sequences have been inserted as indicated below (positions 1 to 15 refer to 3'-terminal nucleotides in the resulting monocistronic vRNAs).

(B) CAT assays as determined relative to pHL1844 (monocistronic CAT construct) after DNA transfection of 293T cells plus helper virus infection followed by one round of progeny viral propagation on MDCK cells are indicated below the lanes. Relative activities of the internal promoter sequences as indicated above the figure refer to measurements in a monocistronic external location of the same promoter variants (Flick *et al.*, Promoter elements in the influenza vRNA terminal structure, RNA 2, 1046-1057 (1996)). Control clone pHL2300 contains an unrelated, non-functional sequence in the central location in an otherwise identical plasmid construct.

Fig. 3. Tandem bicistronic vRNA sup-orting an alternative mode of tran-scription and replication initiation:

An additional internal 3' promoter sequence has been inserted in between both cistrons, in a vRNA-central position. Left half: bicistronic replication and transcription leading to (proximal) GFP expression. Right half: internal initiation resulting in monocistronic replication and transcription leading to (distal) CAT expression, and causing deletion of the GFP sequence from progeny molecules:

Fig. 4. Outgrowth of promoter-up recombinant vRNA versus wildtype vRNA segments in stepwise propagation of influenza virus:

RNA polymerase I transcription of transfected pHL2969 DNA results in influenza vRNA carrying (an external) promoter-up mutant '1104', and containing 2x2 ribozyme targets in flanking positions relative to its HA coding sequence. Another HA vRNA segment in wild-type configuration and originating from infecting FPV helper virus is also present in the recombinant virus preparation, initially (lane 1; 293T lysate) in surplus amounts, but reduced and finally lost entirely in consecutive steps of propagation (lanes 2 to 4; MDCK cell lysates), and in isolated strains after pAM403 ribozyme treatment for removal of the external '1104' promoter sequence (lane 5). Determination throughout by RT-PCR analyses using a pair of primers extending across the 5' inserted target site sequence, with 435 bp representing the recombinant HA segment, and 306 bp the wild-type sequence without an inserted target site sequence.

Fig.5. pAM403 ribozyme cleavage of pHL2969 derived vRNA molecules at specific target sites inserted between an external and an internal 3' promoter sequence:

The external promoter-up ('1104') signal is used for vRNA amplification within recombinant viruses and reduction of helper virus HA vRNA (Fig. 4), while the 'switch' to an internal wild-type signal guarantees stable replication of recombinant viruses. pAM403 hammerhead ribozyme RNAs are indicated in complementary binding to their target site sequences (12 and 10 nucleotides flanking the GU'C cleavage point) by straight lines flanking a central secondary structure symbol. vRNA-internal 2x2 ribozyme targets are marked by xx (see Fig. 7).

Fig. 6. Comparative cleavage analysis of model CAT vRNAs with tandem target sites in various flanking positions, by target-specific ribozymes:

293T tissue culture cells have been transiently DNA-transfected either by a single-headed hammerhead ribozyme (s), or a double-headed (d), or triple-headed (t) ribozyme cDNA construct, all specifically designed to hybridize to a tandem dimer target site sequence inserted in flanking positions into the CAT vRNA. All ribozyme RNAs have been expressed from the same pSV2-*neo* plasmid vector, including a pSV2-*neo* control construct without an inserted ribozyme cDNA sequence (c). At 20 h after DNA transfection (which reached 65% yield as measured by p_{CMV} -GFP transfection in parallel of the same cell culture) the 293T cells were infected by CAT recombinant viruses carrying tandem double target sequences either only in vRNA-3' position, or in both vRNA-3' and 5' positions, or in both vRNA-3' and cRNA-3' positions. Most effective among the s, d, or t-ribozymes were double-headed constructs, acting on 2x2 targets inserted in either of the two localizations described (lanes 6 and 10).

Fig. 7. Alignment of pAM424 double-headed ribozyme with one of their repetitive target sequences located within the 5' and 3' vRNA flanking regions:

The superior activity of ribozymes oriented against targets located in the 3' end of vRNA molecules over those present in the 5' end instead (not shown) is in agreement with the model for influenza vRNA transcription and replication (Lamb and Krug, Orthomyxoviridae: The viruses and their replication. In 'Virology' (B.N.Fields, D.M. Knipe, P.M.Howley, R.M.Chanock, J.L.Melnick, T.P.Monath, B.Roizman, and S.E.Straus, Eds.), 3rd ed., Vol. 1, pp. 1353-1395. Lippincott-Raven, Philadelphia (1996)), according to which influenza polymerase stays attached to the 5' end of the vRNA molecule throughout the entire or even several rounds of transcription, whereas the very 3' end repeatedly, in every initiation reaction serves as the template sequence, and consequently is no longer covered by polymerase. Superiority of a double-headed over a single-headed ribozyme has been determined earlier in this laboratory (A.Menke, Anti-Influenza Ribozyme: vRNA-Spaltung und intrazelluläre Aktivität. Dissertation Universität Giessen (1997)), but the substantial increase of vRNA inactivation rates upon incorporation of tandem target sites at both ends of the vRNA molecule instead of only one has been observed here for the first time, within that overall design.

Fig. 8: pAM424 ribozyme cleavage of resistant FPV wild-type HA vRNA and ribozyme-sensitive pHL2969-derived HA-vRNA in 293T cells infected by vHM41 after isolation from pHL2969-recombinant viral preparations. Lane 1: FPV infection of 293T cells, untreated; lane 2: FPV infection of 293T cells DNA-transfected by pAM424; lane 3: vHM41 infection of 293T cells, untreated; lane 4: vHM41 infection of 293T cells DNA-transfected by pAM424. RT-PCR analyses of purified viral progeny as in Fig. 4.

Fig. 9 Functional analysis of the influenza cRNA promoter structure:

(A) Schematic cRNA promoter ('1104') secondary structure according to the 'corkscrew' model; nucleotides involved in single or double nucleotide exchange are marked by their position.
(B) CAT analyses of 293T cell lysates after DNA transfection and FPV helper virus infection of cRNA promoter variants, in comparison to standard vRNA promoter-up mutant '1104' (pHL1844). Nucleotide substitutions divergent from the basic '1104' structure as present in pHL2583 or pHL2721 (see above) are indicated above the lanes, positions 3 or 8 as marked by a bar refer to cRNA positions counted from the 3' end. Relative CAT activities are marked below the lanes.

Fig. 10 Functional analysis of the vRNA and cRNA promoter in ambisense arrangement:

(A) Sequence organisation of the ambisense promoter cDNA construct carrying T₆/A₆ elements adjacent to the terminal sequence, and secondary structure predictions for the resulting cRNA and vRNA promoter signal.
(B) CAT expression data obtained from the cell lysates of 293T cell after plasmid DNA transfection and FPV infection, and (C) from cell lysates of MDCK cells after one step of viral passage. Indicated above the lanes are promoter/gene conjunctions: v = vRNA promoter; c = cRNA promoter.

Fig.11 Basepair substitutions according to the vRNA 'corkscrew' structure:

(A) 'Corkscrew' conformation of the vRNA promoter drawn against a schematic indication of interacting tripartite viral polymerase. Paired positions exchanged in individual experiments are indicated by numbers, nucleotides 3 or 8 are counted from the 3' end. pHL2024 containing promoter-up mutation '1104' is used as the reference construct (=100%) in all of the CAT assays, while pHL2428 represents the wild-type promoter structure.
(B) CAT expression data obtained after one step of viral passage in MDCK undiluted, and 50 fold diluted.

Fig. 12: Flow-chart of the isolation procedure for an ambisense recombinant influenza virus.

Fig. 13 Immuno-electron microscopy of purified influenza FPV/CSFV-E2-HA virions: Recombinant viruses exposing the foreign glycoprotein CSFV-E2 in their envelopes, which has been fused onto the HA anchor domain, are marked by anti-E2 monospecific antibody and by secondary gold-labelled (5nm) goat antibody. Recombinant viruses (16%) are present together with their FPV helper viruses.

Fig. 14: pHL2969; the exact sequence of the 4930 bps circular DNA is shown in SEQ.ID NO:1.

Fig. 15: pAM403; the exact sequence of the 5811 bps circular DNA is shown in SEQ. ID NO:2.

Fig. 16: pAM424; the exact sequence of the 5860 bps circular DNA is shown in SEQ. ID NO:5.

Fig. 17: pHL2507; the exact sequence of the 4610 bps circular DNA is shown in SEQ. ID NO:6.

Fig. 18: pHL2583; the exact sequence of the 3558 bps circular DNA is shown in SEQ. ID NO:7.

Fig. 19: pHL2989; the exact sequence of the 4343 bps circular DNA is shown in SEQ. ID NO:8.

Fig. 20: pHL1920; the exact sequence of the 3888 bps circular DNA is shown in SEQ. ID NO:9.

[0046] Hence, the present invention is based on two surprising findings, namely

1. influenza virus promoters are active when present internally in a gene;
2. the so-called cRNA, thought to be an intermediate in replication can be turned into a protein-encoding RNA by equipping it with a variant influenza virus promoter, described in the present invention.

[0047] These two observations were used to make ambisense constructs. This allows to package an additional, foreign gene into influenza virus particles. Such particles were made previously, by other methods, but proved to be unstable, and therefore useless. For use as a vaccine for example, a helper virus would have been needed as a stabilizer. Stabilization in the present invention is achieved by several means. These include the "balancing" of one of the two promoters in the ambisense bicistronic genetic construct with seven other vRNA wildtype promoters, while the additional promoter is used for high-rate expression of the foreign gene at various levels.

[0048] Thus, the present invention provides a system for expression of foreign proteins in higher eukaryotic systems. One system in particular is interesting, namely embryonated chicken eggs, as it allows cost-effective production in an automatable way (as used by most flu vaccine producers). The reason that this process is now possible, is that the foreign protein is part of a stable, engineered influenza virus particle. The virus can be designed also to rapidly monitor process improvements.

[0049] An excellent use is of course the use of the construct as a vaccine. The influenza virus particle is immunogenic and can now be equipped with foreign antigens, enabling for example the design and production of hepatitis C virus and HIV vaccines, but also of tumor vaccines. As the present invention shows, the foreign antigenic surface glycoprotein is "fused" to a C-terminal segment of influenza HA, and the antigen then is presented at the surface of influenza virus particles. In addition, these vaccines can now be made in the way standard flu vaccines are made, i.e., in embryonated chicken eggs.

SEQUENCE LISTING

<110> ARTEMIS PHARMACEUTICALS GmbH

<120> Stable Recombinant Influenza Viruses Free of Helper
Viruses

<130> 990383ep

<140>

<141>

<160> 9

<170> PatentIn Ver. 2.1

<210> 1

<211> 4930

<212> DNA

<213> Influenza virus

<400> 1

```

cgtacgaagc ttctagaggg attggctgag acgaaaaaca tatgctagag ggattggctg 60
agacgaaaaa catatgctag agcggccgcc accgcggtgg agctccagct tttgttcctt 120
25 ttagttaggg ttaattgcgc gcaggcctag ctaggtaaag aaaaataccc ttgtttctac 180
taataaccgc gcgccccaaa atgcgcgactc ggagcgaaag atatacctcc cccggggcgcg 240
ggaggtcgcg tcaccgacca cgccgcgcgc ccaggcgacg cgcgacacgg acacctgtcc 300
ccaaaaacgc caccatcgca gccacacacg gaggcgcccg ggccctcttg lcaacccag 360
gacacacgcg ggagcagcgc cgggcccggg acgcccctcc ggccgcccgt gccacacgca 420
30 gggggcccgcc ccgtgtctcc agagcgggag ccggaagcat ttccggcccg cccctctac 480
gaccgggaca cagagggac cgaaggccgc ccaggcgca cctctcgcc cgcacgcgcg 540
ctcagggagc gctctccgac tccgcacggg gactcgccag aaaggatcgt gacctgcatt 600
aatgaatcag gggataacgc aggaaagaac atgtgagcaa aaggccagca aaaggccagg 660
aaccgtaaaa aggcgcgctt gctggcgctt ttccataggc tccgcccccc tgacgagcat 720
35 cacaaaaatc gacgctcaag tcagaggtgg cgaaacccga caggactata aagataccag 780
gcggttcccc ctggaagctc cctcgtgcgc tctcctgttc cgacctgcc gcttaccgga 840
tacctgtccg cctttctccc ttcggaagc gtggcgcttt ctcatagctc acgctgtagg 900
tatctcagtt cgggtgtaggt cgttcgctcc aagctggggt gtgtgcacga acccccgtt 960
cagcccgacc gctgcgcctt atccggtaac tategtcttg agtccaaccc ggtaagacac 1020
40 gacttatcgc cactggcagc agccactggt aacaggatta gcagagcgag gtatgtaggc 1080
gggtctacag agttcttgaa gtggtggcct aactacggct acactagaag gacagtattt 1140
gggtatctcg ctctgctgaa gccagttacc ttcggaaaaa gagttagtag ctcttgatcc 1200
ggcaaacaaa ccaccgctgg tagcggtggt tttttgttt gcaagcagca gattacgcgc 1260
agaaaaaaag gatctcaaga agatcccttg atcttttcta cggggtctga cgctcagtg 1320
45 aacgaaaact caggttaagg gatcttggtc atgagattat caaaaaggat cttcacctag 1380
atccttttaa attaaaaatg aagtttttaa tcaatctaaa gtatatatga gtaaaccttg 1440
tctgacagtt accaatgctt aatcagtgag gcacctatct cagcgatctg tctatttcgt 1500
tcatccatag ttgcctgact cccgctcgtg tagataacta cgatacggga gggcttacca 1560
tctggcccca gtgctgcaat gataccgcga gaccacgct caccggctcc agatttatca 1620
50 gcaataaaac agccagcccg aagggccgag cgcagaagtg gtcttgcaac tttatccgcc 1680
tccatccagt ctattaattg ttgcccggaa gctagagtaa gtagttcgcc agttaatagt 1740
ttgcgcaacg ttgttgccat tgctacagcg atcgtggtgt cagctcgtc gtttggtatg 1800
gcttcattca gctccggttc ccaacgatca aggcagttc catgatcccc catgttggtc 1860
aaaaaagcgg ttagctcctt cgggtctccg atcgttgtca gaagtaagtt ggccgcatg 1920
55 ttatcactca tggttatggc agcactgcat aattctctta ctgcatgcc atccgtaaga 1980

```

5 tgcttttctg tgactggtga gtactcaacc aagtcattct gagaatagtg tatgcggcga 2040
 ccgagttgct cttgcccggc gtcaacacgg gataataccg cgccacatag cagaacttta 2100
 aaagtgtca tcattgaaa acgttcttcg gggcgaaaac tctcaaggat cttaccgctg 2160
 ttgagatcca gttcgatgta acccactcgt gcacccaact gatcttcagc atcttttact 2220
 ttcaccagcg tttctgggtg agcaaaaaa ggaaggcaaa atgccgcaaa aaaggggaata 2280
 agggcgacac ggaaatgttg aatactcata ctcttccttt ttcaatatta tgaagcatt 2340
 tatcaggggt attgtctcat gagcggatac atatttgaat gtatttagaa aaataaacaa 2400
 aagagtttgt agaaacgcaa aaaggccatc cgtcaggatg gccttctgct taatttgatg 2460
 10 cctggcagtt tatggcgggc gtctgccccg ccacccctcg ggccgttgct tcgcaacgtt 2520
 caaatccgct cccggcggat ttgtcttact caggagagcg ttcaccgaca aacaacagat 2580
 aaaacgaaag gccagtcctt tcgactgagc ctttcgtttt atttgatgcc tggcagttcc 2640
 ctactctcgc atggggagac cccacactac categgcgct acggcgtttc acttctgagt 2700
 tcggcatggg gtcaggtggg accaccgcgc tactgccgcc aggcaaatc tgttttatca 2760
 15 gaccgcttct gcgttctgat ttaatctgta tcaggctgaa aatcttctct catccgcca 2820
 aacagaagct agcggccgat ccccaaaaaa aaaaaaaaaa aaaaaaaaaa gagtccagag 2880
 tggccccgcc gttccgcgcc gggggggggg gggggggggg acacttctcg acatctggtc 2940
 gacctccagc atcgggggaa aaaaaaaaaa caaagtctcg cccggagtac tggtcgacct 3000
 ccgaagtggg gggggagtag aaacagggtg gataatcact cactgacgta cgttgagcaa 3060
 20 ctgactgaaa tgccttgagc aactgactga aatgcctgac gtctttagca aaagcagggt 3120
 agataatcac tctgtagtg acatccacat cgtaccagga ttggctgaga cgaaaaacat 3180
 attgtaccag ggtttggctg agacgaaaaa cctattgtag gtacaaaaat gaacactcaa 3240
 atcttggttt tcgccttgc ggtagtcac cccacaaatg cagacaaaat ttgtcttggg 3300
 catcatgctg tatcaaatgg caccaaagta aacacactca ctgagagagg agtagaagtt 3360
 25 gtcaatgcaa cggaaacagt ggagcggaca aacatcccca aaatttgctc aaaagggaaa 3420
 agaaccactg atcttgacca atgcggactg ttagggacca ttaccggacc acctcaatgc 3480
 gaccaatttc tagaattttc agctgatcta ataatcgaga gacgagaagg aaatgatgtt 3540
 tgttaccctg ggaagtgtgt taatgaagag gcattgcgac aaatcctcag aggatcaggt 3600
 gggattgaca aagaaacaat gggattcaca tatagtggaa taaggaccaa cggaacaact 3660
 30 agtgcagtga gaagatcagg gtcttcattc tatgcagaaa tggagtggct cctgtcaaat 3720
 acagacaatg cttctttccc acaaatgaca aaatcataca aaaacacagg gagagaatca 3780
 gctctgatag tctggggaat ccaccattca ggatcaacca ccgaacagac caaactatat 3840
 gggagtggaa ataaactgat aacagtcggg agttccaaat atcatcaatc tttgtgcccg 3900
 agtccaggaa cacgaccgca gataaatggc cggtccggac ggattgattt tcattgggtg 3960
 atcttggatc ccaatgatac agttactttt agtttcaatg gggctttcat agctccaaat 4020
 35 cgtgccagct tcttgagggg aaagtccatg gggatccaga gcgatgtgca ggttgatgct 4080
 aattgcgaag gggaatgcta ccacagtggg gggactataa caagcagatt gccttttcaa 4140
 aacataaata gcagagcagt tggcaaatgc ccaagatatg taaaacagga aagtttatta 4200
 ttggcaactg ggatgaagaa cgttcccga ccttccaaaa aaaggaaaaa aagaggcctg 4260
 tttggtgcta tagcaggggt tattgaaaat ggttgggaag gtctgggtcg cgggtgggtac 4320
 40 ggtttcaggc atcagaatgc acaaggagaa ggaactgcag cagactacaa aagcacccaa 4380
 tcggcaattg atcagataac cggaaggtta aatagactca ttaagaaaac caaccagcaa 4440
 tttgagctaa tagataatga attcactgaa gtggaaaagc agattggcaa ttttaattaac 4500
 tggaccaaag actccatcac agaagtatgg tcttacaatg ctgaacttct tgtggcaatg 4560
 45 gaaaaccagc acactattga tttggctgat tcagagatga acaagctgta tgagcgagtg 4620
 aggaaacaat taagggaaaa tgctgaagag gatggcactg gttgctttga aatttttcat 4680
 aaatgtgacg atgattgtat ggctagtata aggaacaata cttatgatca cagcaaatat 4740
 agagaagaag cgatgcaaaa tagaatacaa attgaccag tcaaattgag tagtggctac 4800
 aaagatgtga tactttgggt tagcttcggg gcacatgct ttttgcttct tgccattgca 4860
 atgggccttg ttttcatatg tgtgaagaac ggaaacatgc ggtgcactat atgcatttaa 4920
 50 agcttgcatg 4930

<210> 2

<211> 5811

<212> DNA

55 <213> Influenza virus

<400> 2

aattcctttg cclaaallaa atgaggactt aacctgtgga aatattttga tgtgggaagc 60
 5 tgttactgtt aaaactgagg ttattggggg aactgctatg ttaaacttgc attcagggac 120
 acaaaaaact catgaaaatg gtgctggaaa acccattcaa gggccaaatt ttcatttttt 180
 tgctgttggt ggggaacctt tggagctgca ggtgtgttta gcaactaca ggaaccaata 240
 tcctgctcaa actgtaaccc caaaaaatgc tacagttgac agtcagcaga tgaacactga 300
 ccacaaggct gttttggata aggataatgc ttatccagtg gagtgctggg ttcctgatcc 360
 10 aagtaaaaaat gaaaacacta gatatttttg aacctacaca ggtggggaaa atgtgcctcc 420
 tgttttgcac attactaaca cagcaaccac agtgcctctt gatgagcagg gtgttggggc 480
 cttgtgcaaa gctgacagct tgtatgtttc tgctgttgac atttgtgggc tgtttacca 540
 cacttctgga acacagcagt ggaagggact tcccagatat tttaaaatta cccttagaaa 600
 gcggtctgtg aaaaacccct acccaatttc ctttttgtta agtgacctaa ttaacaggag 660
 15 gacacagagg gtggatgggc agcctatgat tggaatgtcc tctcaagtag aggagggttag 720
 ggtttatgag gacacagagg agcttcctgg ggatccagac atgataagat acattgatga 780
 gtttggaaca accacaacta gaatgcagtg aaaaaaatgc tttatttctg aaatttctga 840
 tgctattgct ttatttctaa ccattataag ctgcaataaa caagttaaca acaacaattg 900
 cattcatttt atgtttcagg ttcaggggga ggtgtgggag gttttttaa gcaagtaaaa 960
 20 cctctacaaa tgtggtatgg ctgattatga tctctagtca aggcactata catcaaatat 1020
 tccttattaa cccctttaca aattaaaaag ctaaagggtac acaatttttg agcatagtta 1080
 ttaatagcag aactctatg cctgtgtgga gtaagaaaaa acagtatgtt atgattataa 1140
 ctgttatgcc tacttataaa gggtacagaa ttttttcca taattttctt gtatagcagt 1200
 gcagcttttt cctttgtggt gtaaataagca aagcaagcaa gagttctatt actaaacaca 1260
 25 gcatgactca aaaaacttag caattctgaa ggaaagtcct tggggtcttc tacctttctc 1320
 ttcttttttg gaggagtaga atgttgagag tcagcagtag cctcatcctc actagatggc 1380
 atttcttctg agcaaaacag gttttcctca ttaaaggcat tccaccactg ctcccatcca 1440
 tcagttccat agyllgyaal claaaatata caaacaatta gaatcagtag tttaacacat 1500
 tatacctta aaaattttat atttacctta gagctttaaa tctctgtagg tagtttctcc 1560
 aattatgtca caccacagaa gtaaggttcc ttcacaaaga tccgggacca aagcggccat 1620
 30 cgtgcctccc cactcctgca gttcgggggc atggatgcgc ggatagccgc tgctgggtttc 1680
 ctggatgccg acggatttgc actgccggta gaactccgcg aggtcgtcca gcctcaggca 1740
 gcagctgaac caactcgcga ggggatcgag cccggggtgg gcgaagaact ccagcatgag 1800
 atccccgcgc tggaggatca tccagccggc gtcccgaaa acgattccga agcccaacct 1860
 35 ttcataagaag gcggcggttg aatcgaaatc tcgtgatggc aggttgggag tcgcttggtc 1920
 ggtcatttcg atgaattcga gctcggtagc cggggatcct ctagaggcat ttcagtttcg 1980
 tcctcacgga ctcatcagag ttgctcaatt cgaaccccag agtcccgcctc agaagaactc 2040
 gtcaagaagg cgatagaagg cgatgcgtg cgaatcggga gcggcgatc cgtaaagcac 2100
 gaggaagcgg tcagcccatc cgcgcgcaag ctcttcagca atatcacggg tagccaacgc 2160
 40 tatgtcctga tagcgggtccg ccacaccag cgggccacag tcgatgaatc cagaaaagcg 2220
 gccattttcc accatgatat tcgggaagca ggcacgcga tgggtcacga cgagatcctc 2280
 gccgtcgggc atgcgcgcct tgagcctggc gaacagttcg gctggcgcca gccctgatg 2340
 ctcttcgtcc agatcatcct gatcgacaag accggcttcc atccgagtag gtgctcgctc 2400
 gatgcgatgt ttcgcttggt ggtcgaaatg gcaggtagcc ggatcaagcg tatgcagccg 2460
 ccgcattgca tcagccatga tggatacttt ctggcagga gcaaggtgag atgacaggag 2520
 45 atcctgcccc ggcacttcgc ccaatagcag ccagtccctt cccgcttcag tgacaacgct 2580
 gagcacagct gcgcaaggaa cgcccgtcgt ggccagccac gatagccgcg ctgcctcgctc 2640
 ctgcagttca ttcagggcac cggacagggtc ggtcttgaca aaaagaaccg ggcgcccctg 2700
 cgctgacagc cggaacacgg cggcatcaga gcagccgatt gtctgttgtg cccagtcata 2760
 gccgaatagc ctctccaccc aagcggccgg agaacctgcg tgcaatccat cttgttcaat 2820
 50 catgcaaaac gatccctac tctgtctctt atcagatcct gatccccgc gccatcagat 2880
 ccttggcggc aagaaagcca tccagtttac tttgcaggyc tccccacct taccagaggg 2940
 cgccccagct ggcaattccg gttcgcttgc tgtccataaa accgcccagt ctagctatcg 3000
 ccatgtaagc ccaactgcaag ctacctgctt tctctttgcg cttgcgtttt cccttgcca 3060
 gatagcccag tagctgacat tcatccgggg tcagcacctt tctgcggac tggctttcta 3120
 55 cgtgttccgc ttccttttag agcccttgcg cctgagtgcc ttgcggcagc gtgaagcttt 3180

ttgcaaaaagc ctaggcctcc aaaaaagcct cctcactact tctggaaag ctcagaggcc 3240
 gaggcggcct cggcctctgc ataaataaaa aaaattagtc agccatgggg cggagaatgg 3300
 gcggaactgg gcggagttag gggcgggatg ggcggagtta ggggcyygac latgggtgct 3360
 gactaattga gatgcatgct ttgcatactt ctgcctgctg gggagcctgg ggactttcca 3420
 cacctgggtg ctgactaatt gagatgcatg ctttgcatac ttctgcctgc tggggagcct 3480
 ggggactttc cacaccctaa ctgacacaca ttccacagct gcctcgcgcg tttcggtgat 3540
 gacggtgaaa acctctgaca catgcagctc ccggagacgg tcacagcttg tctgtaagcg 3600
 gatgcgggga gcagacaagc ccgtcagggc gcgtcagcgg gtgttggcgg gtgtcggggc 3660
 gcagccatga cccagtcacg tagcgatagc ggagtgtata ctggcttaac tatgcggcat 3720
 cagagcagat tgtactgaga gtgcaccata tgcgggtgtga aataccgcac agatgcgtaa 3780
 ggagaaaaata ccgcctcagg cgctcttcgg cttctcggct cactgactcg ctgcgctcgg 3840
 tcgttcggct gcggcgagcg gtatcagctc actcaaaggc ggtaatacgg ttatccacag 3900
 aatcagggga taacgcagga aagaacatgt gagcaaaagg ccagcaaaag gccaggaacc 3960
 gtaaaaaggc cgcgttgctg gcgtttttcc ataggctcgg ccccccctgac gagcatcaca 4020
 aaaaatcgacg ctcaagtcag aggtggcgaa acccgacagg actataaaga taccaggcgt 4080
 tttccccttg aaagctccctc gtgcgctctc ctgttccgac cctgccgctt accggatacc 4140
 tgtccgcctt tctcccttcg ggaagcgtgg cgctttctca tagctcacgc tgtaggtatc 4200
 tcagttcggg taggtcggt cgctccaagc tgggctgtgt gcacgaacct cccgttcagc 4260
 ccgaccgctg cgccttatcc ggtaactatc gtcttgagtc caaccggta agacacgact 4320
 tatcgccact ggcagcagcc actggttaaca ggattagcag agcgaggat gtaggcgggtg 4380
 ctacagagtt cttgaagtgg tggcctaact acggctacac tagaaggaca gtattttggtg 4440
 tctgcgctct gctgaagcca gttaccttcg gaaaaagagt tggtagctct tgatccggca 4500
 aacaaaccac cgctggtagc ggtggttttt ttgtttgcaa gcagcagatt acgcgcagaa 4560
 aaaaaggatc tcaagaagat cctttgatct tttctacggg gtctgacgct cagtggaaacg 4620
 aaaaactcacg ttaagggtatt ttggtcatga gattatcaaa aaggatcttc acctagatcc 4680
 ttttaaattg aaaaatgaagt tttaaatcaa tctaaagtat atatgagtaa acttgggtctg 4740
 acagttacca atgcttaac agtgaggcac ctatctcagc gatctgtcta tttcgttcat 4800
 ccatagttgc ctgactcccc gtcgtgtaga taactacgat acgggagggc ttaccatctg 4860
 gccccagtg tgcaatgata ccgcgagacc cacgctcacc ggctccagat ttatcagcaa 4920
 taaaccagcc agccggaagg gcgagcgca gaagtgttcc tgcaacttta tccgcctcca 4980
 tccagttctat taattgttgc cgggaagcta gagtaagtag ttcgccagtt aatagtttgc 5040
 gcaacgttgt tgccattgct gcaggcatcg tgggtgcacg ctcgctcgctt ggtatggctt 5100
 cattcagctc cggttcccaa cgatcaaggc gagttacatg atcccccatg ttgtgcaaaa 5160
 aagcggttag ctcttcgggt cctccgatcg ttgtcagaag taagttggcc gcagtgttat 5220
 cactcatggt tatggcagca ctgcataatt ctcttactgt catgccatcc gtaagatgct 5280
 tttctgtgac tgggtgagta tcaaccaagt cattctgaga atagtgtatg cggcgaccga 5340
 gttgtctctg cccggcgctc acacgggata ataccgcgcc acatagcaga actttaaaag 5400
 tgctcatcat tggaaaacgt tcttcggggc gaaaactctc aaggatctta ccgctgttga 5460
 gatccagttc gatgtaacct actcgtgcac ccaactgatc ttcagcatct tttactttca 5520
 ccagcgtttc tgggtgagca aaaacaggaa ggcaaaatgc cgcaaaaaag ggaataaggg 5580
 cgacacggaa atggtgaata ctacatactt tcccttttca atattattga agcattttatc 5640
 agggttattg tctcatgagc ggatacatat ttgaatgtat ttagaaaaat aaacaaatag 5700
 gggttccgcg cacatttccc cgaaaagtgc cacctgacgt ctaagaaacc attattatca 5760
 tgacattaac ctataaaaaat aggcgtatca cgaggccctt tcgtcttcaa g 5811

<210> 3

<211> 2005

<212> DNA

<213> Influenza virus

<400> 3

agtagaaaca aggggtatatt tctttacctg gctaggcctg cgcgcaatta accctcacta 60
 aagggaaaca aagctggagc tccaccgcgg tggcgccgc tctagcatat gtttttcgtc 120
 tcagccaatc cctctagcat atgtttttcg tctcagccaa tccctctaga agcttcgtac 180
 gcatgcaagc tttaaatgca tatagtgcac cgcattgttc cgttcttcac acatatgaaa 240

acaaggccca ttgcaatggc aagaagcaaa aagcatgatg ccccgaaagt aaaccaaagt 300
 atcacatctt ttagccact actcaatttg actgggtcaa tttgtattcl attttgatc 360
 gcttcttctc tgtatttgct gtgatcataa gtattgttcc ttatactagc catacaatca 420
 5 tegt.cacatt tatgaaaaat ttcaaagcaa ccagtgccat cctcttcagc attttccctt 480
 aattgttttc tcaactcgctc atacagcttg ttcatctctg aatcagccaa atcaatagt 540
 tgetgggttt ccattgccac aagaagtcca gcattgtaag accatacttc tgtgatggag 600
 tctttgggtc agttaattaa attgccaatc tgcttttcca cttcagtga ttcatatct 660
 attagctcaa attgctgggt gggttttctta atgagtctat ttaactttcc ggttatctga 720
 tcaattgccg attgggtgct tttgtagtct gctgcagttc cttctccttg tgcattctga 780
 10 tgcctgaaac cgtaccaccc gtgcaccaga ctttcccaac cattttcaat aaacctgct 840
 atagcaccaa acaggcctct ttttttctt tttttggaag gttcgggaac gttcttcac 900
 ccagttgcca ataataaact ttctgtttt acatatcttg ggcatttgcc aactgctctg 960
 ctatttatgt tttgaaaagg caatctgctt gttatagtcc ctccactgtg gtagcattcc 1020
 ccttcgcaat tagcatcaac ctgcacatcg ctctggatcc ccattggactt tcccccaag 1080
 15 aagctggcac gatttgagc tatgaaagcc ccattgaaac taaaagtaac tgtatcattg 1140
 ggatccaaga tcaaccaatg aaaatcaatc cgtccggacc ggccatttat ctgcggtcgt 1200
 gttcctggac tgcgcacaaa agattgatga tatttggaac tcccgaactg tatcagttta 1260
 tttccactcc catatagttt ggtctgttcg gtggttgatc ctgaatggtg gattccccag 1320
 actatcagag ctgattctct cctgtgttt ttgtatgatt ttgtcatttg tgggaaagaa 1380
 20 gcattgtctg tatttgacag gagccactcc atttctgcat agaataaga cctgatctt 1440
 ctacatgcac tagttgttcc gttggtcctt attccactat atgtgaatcc cattgtttct 1500
 ttgtcaatcc cacctgatcc tctgaggatt tgtcgcaatg cctcttcatt aacaaacttc 1560
 cccgggtaac aaacatcatt tcttctcgt ctctcgatta ttagatcagc tgaaaattct 1620
 agaaattggt cgcattgagg tggccggta atggcccta acagtcgca ttggccaaga 1680
 25 tcagtgggtc ttttcccttt tgagcaaat ttggggatgt ttgtccgctc cactgtttcc 1740
 gttgcattga caacttctac tctctctca gtgagtgtgt ttactttggt gccatttgat 1800
 acagcatgat gtccaagaca aattttgtct gcatttggtg ggatgactgc cgcaaggcg 1860
 aaaaccagga tttgagtgtt cattttggtt cctacaatat gttttctgct tcagccaatc 1920
 cctggtacaa tatgtttttc gtctcagcca atctcgttac gatgtggatg tcaactcagt 1980
 30 agtgattatc taccctgctt ttgct 2005

<210> 4

<211> 1146

<212> DNA

35 <213> Influenza virus

<400> 4

agtagaaaca aggtatattt tctttaccta gctaggcctg cgcgcaatta accctcacta 60
 aagggaaaca aagctggagc tccaccgcgg tggcgccgc tctagcatat gtttttcgtc 120
 40 tcagccaatc cctctagcat atgtttttcg tctcagccaa tccctctaga agcttcgtac 180
 gcatgcttaa ataagctgaa acgagaaagt tcttatctct tgctccactt caagcgtag 240
 ttgtaaggct tgcataaatg ttatttggtc aaaactatcc tctgttatct tcaatctatg 300
 tctcacttct tcaattaacc atcttatctt ttcaaatttc tgactcaatt gttctcgcca 360
 ttttccgttt ctgctttgga gggagtggag gtcccccat ctactactg cttctccaag 420
 45 cgaatctctg tatagtttca gagactcgaa ctgtgttatc attccattca agtctccga 480
 tgaggacccc aattgcattt ttgacatcct catcagtatg tcttggaga gaaggcaatg 540
 gtgaaatttc gccgacaatt gtcctctcat cgggttaaagc ccttaatagt atgagagttt 600
 ccagccgatc gaaaatcaca ctgaagtttg ctttcagtat gatgttcttc cccatgatcg 660
 cctggtccat tctgatgcaa agggagcctg ccactttctg tttgggcatg agcatgaacc 720
 50 agtcccttga catctcttca agagtcatgt cagttaggta gcgtgtagca ggtacagagg 780
 caatggctcat ttttaagtgc tcatcggatt cgtcctccag aatccgctcc actatctgct 840
 ttccaacacg agtagctgtg tcatgttcca gaccaagagt gctgcctctt cccctcagg 900
 acttctgac tccggcaagt cggccaagga atggggcatc acccatttct tgggtctgca 960
 atcgtttgct gacatgccaa agaaagcagt ctacctgaaa gcttgacaca gtgttggaal 1020
 55 ccattatggt acctacaata tgtttttcgt ctcagccaat ccctggtaca atatgttttt 1080

EP 1 035 209 A1

cgtctcagcc aatcctggta cgatgtggat gtcactcagt gagtgattat ctaccctgct 1140
tttgct 1146

<210> 5
<211> 5860
<212> DNA
<213> Influenza virus

<400> 5
catcgattgg ctgactgatg agtccgtgag gacgaaacga aaaacatatt gtagagctcg 60
aattcatcga aatgacggac caagcgacgc ccaacctgcc atcacgagat ttcgattcca 120
cgcgcgctt ctatgaaagg ttgggcttcg gaatcgtttt cggggacgcc ggctggatga 180
tctccagcg cggggatctc atgctggagt tcttcgccc ccccggtctc gatccctcg 240
cgagtgtgtt cagctgctgc ctgaggctgg acgacctcgc ggagttctac cggcagtga 300
aatccgctcg catccaggaa accagcagcg gctatccgcg catccatgcc cccgaactgc 360
aggagtgggg aggacgatg gccgctttgg tcccggatct ttgtgaagga accttacttc 420
tgtgtgtga cataattgga caaactacct acagagattt aaagctctaa ggtaaatata 480
aaatttttaa gtgtataatg tgttaacta ctgattctaa ttgtttgtgt attttagatt 540
ccaacctatg gaactgatga atgggagcag tgggtgaatg cctttaatga ggaaaacctg 600
ttttgtctcag aagaaatgcc atctagtgat gatgaggcta ctgctgactc tcaacattct 660
actcctccaa aaaagaagag aaaggtagaa gacccaagg actttccttc agaattgcta 720
agttttttga gtcatgctgt gtttagtaat agaactcttg cttgctttgc tatttacacc 780
acaaaggaaa agctgcact gctatacaag aaaattatgg aaaaatattc tgtaaccttt 840
ataagtaggc ataacagtta taatcataac atactgtttt ttcttactcc acacaggcat 900
agagtgtctg ctattaataa ctatgctcaa aaattgtgta cctttagctt tttattttgt 960
aaagggttca ataaggaata ttgatgtat agtgccttga ctagagatca taatcagcca 1020
taccacattt gtagagggtt tacttgcctt aaaaaacctc ccacacctcc ccctgaacct 1080
gaaacataaa atgaatgcaa ttgttgttgt taacttgttt attgcagctt ataattggtta 1140
caaataaagc aatagcatca caaatttcac aaataaagca tttttttcac tgcattctag 1200
ttgtgttttg tccaaactca tcaatgtatc ttatcatgtc tggatcccca ggaagctcct 1260
ctgtgtcctc ataaacccta acctcctcta ctgagagga cattccaatc ataggctgcc 1320
catccacctc ctgtgtcctc ctgttaatta ggtcacttaa caaaaaggaa attgggtagg 1380
ggtttttcac agaccgcttt ctaagggtaa ttttaaaata tctgggaagt cccttccact 1440
gctgtgttcc agaagtgttg gtaaacagcc cacaatgtc aacagcagaa acatacaagc 1500
tgtcagcttt gcacaagggc ccaacacctc gctcatcaag aagcactgtg gttgctgtgt 1560
tagtaatgtg caaaacagga ggcacatttt cccacctgt gtaggttcca aaatatctag 1620
tgttttcatt ttacttgcga tcaggaaacc agcactccac tggataagca ttatccttat 1680
ccaaaacagc cttgtgtgca gtgttcatct gctgactgtc aactgtagca ttttttgggg 1740
ttacagtttg agcaggatat ttggctcctgt agtttgcata cacacctgc agctccaaag 1800
gttccccacc aacagcaaaa aaatgaaaat ttgaccttg aatgggtttt ccagcaccat 1860
tttcatgagt tttttgtgtc cctgaatgca agtttaacat agcagttacc ccaataacct 1920
cagttttaac agtaacagct tccacatca aaatatttcc acagggttaag tctcatttta 1980
aattaggcaa aggaattcct gaagacgaaa gggcctcgtg atacgcctat ttttatagg 2040
taatgtcatg ataataatgg tttcttagac gtcagggtggc acttttcggg gaaatgtgcg 2100
cggaaccctt atttgtttat ttttctaaat acattcaaat atgtatccgc tcatgagaca 2160
ataaccctga taaatgttc aataatattg aaaaaggaa agtatgagta ttcaacattt 2220
ccgtgtcgcc cttattccct tttttgcggc attttgcctt cctgtttttg ctacccaga 2280
aacgtgtgtg aaagtataag atgctgaaga tcagttgggt gcacgagtgg gttacatcga 2340
actggatctc aacagcggta agatccttga gaggtttcgc cccgaagaac gttttccaat 2400
gatgagcact tttaaagtgc tgctatgtgg cgcggtatta tcccggtgtg acgcccggca 2460
agagcaactc ggtcgccgca tacactattc tcagaatgac ttggttgagt actcaccagt 2520
cacagaaaag catcttacgg atggcatgac agtaagagaa ttatgcagtg ctgccataac 2580
catgagtgat aacactgcgg ccaacttact tctgacaacg atcggaggac cgaaggagct 2640
aaccgctttt ttgcacaaca tgggggatca tgtaactcgc cttgatcggt gggaaaccgga 2700
gctgaatgaa gccataccaa acgacgagcg tgacaccacg atgcctgcag caatggcaac 2760

aacgttgccg aaactattaa ctggcgaaact acttactcta gcttcccggc aacaattaat 2820
 agactggatg gaggcggata aagttgcagg accacttctg cgttcggccc ttcgggctgg 2880
 ctggtttatt gctgataaat ctggagccgg tgagcgtggg tctcgggta tcattgcagc 2940
 actggggcca gatggtaagc cctcccgtat cgtagttatc tacacgacgg ggagtcaggc 3000
 5 aactatggat gaacgaaata gacagatcgc tgagataggt gcctcactga ttaagcattg 3060
 gtaactgtca gaccaagttt actcatatat acttttagatt gatttaaaac ttcattttta 3120
 atttaaaagg atctaggtga agatcctttt tgataatctc atgacaaaaa tcccttaacg 3180
 tgagttttcg ttcactgag cgtcagaccc cgtagaaaaa atcaaaggat cttcttgaga 3240
 tccttttttt ctgcgcgtaa tctgctgctt gcaaacaaaa aaaccaccgc taccagcggc 3300
 10 ggtttgtttg cgggatcaag agctaccaac tctttttccg aaggtaactg gcttcagcag 3360
 agcgcagata ccaaatactg tccttctagt gtacccgtag ttaggccacc acttcaagaa 3420
 ctctgtagca ccgcctacat acctcgctct gctaatectg ttaccagtgg ctgctgccag 3480
 tggcgataag tcgtgtctta ccgggttgga ctcaagacga tagttaccgg ataaggcgca 3540
 gcggtcgggc tgaacggggg gttcgtgcac acagcccagc ttggagcgaa cgacctacac 3600
 15 cgaactgaga tacctacagc gtgagctatg agaaagcgcc acgcttcccg aaggggagaaa 3660
 ggcggacagg tatccggtaa gcggcagggc cggaaacagga gagcgacga gggagcttcc 3720
 agggggaaac gcctggatc tttatagtc tgtcggggtt cggcacctct gacttgagcg 3780
 tcgatttttg tgatgctcgt cagggggggc gagcctatgg aaaaacgcca gcaacgcggc 3840
 ctttttacgg ttcctggcct tttgctggcc tttgctcac atgttcttcc ctgcgttacc 3900
 20 cctgattct gtgataaac gtattaccgc ctttgagtga gctgataccg ctgcgcgagc 3960
 ccgaacgacc gagcgacgc agtcagttag cgaggaaagc gaagagcgcc tgatgcggta 4020
 ttttctcctt acgcatctgt gcggtatttc acaccgcata tggtgcactc tcagtacaat 4080
 ctgctctgat gccgcatagt taagccagta tacactccgc tatcgctacg tgactgggtc 4140
 atggctgcgc ccgcacaccc gccaacaccc gctgacgcgc cctgacgggc ttgtctgctc 4200
 25 ccggcatccg cttacagaca agctgtgacc gtctccggga gctgcatgtg tcagagggtt 4260
 tcaccgtcat caccgaaacg cgcgaggcag ctgtggaatg tgtgtcagtt aggggtgtgga 4320
 aagtcgccag gctccccagc aggcagaagt atgcaaagca tgcatctcaa ttagtcagca 4380
 accaggtgtg gaaagtcccc aggtccccca gcaggcagaa gtatgcaaag catgcatctc 4440
 aattagtcag caaccatagt cccgccccta actccgccc tcccgcacct aactccgccc 4500
 30 agttccgccc attctccgcc ccatggctga ctaattttt ttattttatgc agaggccgag 4560
 gccgcctcgg cctctcagc attccagaag tactgaggag gcttttttgg aggcctaggc 4620
 ttttgcaaaa agcttcacgc tgccgcaagc actcagggcg caagggctgc taaaggaagc 4680
 ggaacacgta gaaagccagt ccgcagaaac ggtgctgacc ccggatgaat gtcagctact 4740
 gggctatctg gacaagggaa aacgcaagcg caaagagaaa gcaggtagct tgcagtgggc 4800
 35 ttacatggcg atagctagac tgggcgggtt tatggacagc aagcgaaccg gaattgccag 4860
 ctggggcgcc ctctggttaag gttgggaagc cctgcaaagt aaactggatg gctttcttgc 4920
 cgccaaggat ctgatggcgc aggggatcaa gatctgatca agagacagga tgaggatcgt 4980
 ttcgcatgat tgaacaagat ggattgcacg caggttctcc ggccgcttgg gtggagaggc 5040
 tattcggtca tgactgggca caacagacaa tcggtgctc tgatgccgc gtgttccggc 5100
 40 tgtcagcgca ggggcgccc gttctttttg tcaagaccga cctgtccggc gccctgaatg 5160
 aactgcagga cgaggcagcg cggctatcgt ggctggccac gacggcggtt ccttgccgag 5220
 ctgtgctcga cgttgtcact gaagcgggaa gggactggc gctattgggc gaagtgccgg 5280
 ggcaggatct cctgtcatct caccttgctc ctgccgagaa agtatccatc atggctgatg 5340
 caatgcggcg gctgcatacg cttgatccgg ctacctgcc attcgaccac caagcgaaac 5400
 45 atcgcatcga gcgagcacgt actcggatgg aagccggtct tgtcgatcag gatgatctgg 5460
 acgaagagca tcaggggctc gcgccagccg aactgttcgc caggctcaag gcgcgcatgc 5520
 ccgacggcga ggatctcgtc gtgacccatg gcgatgcctg cttgccgaat atcatggtgg 5580
 aaaatggccg cttttctgga ttcacgact gtggccggct ggggtgtggc gaccgctatc 5640
 50 aggacatagc gttggctacc cgtgatattg ctgaagagct tggcggcgaa tgggctgacc 5700
 gcttctcgt gctttacgg atcgcgcgtc ccgattcgca gcgcatcgcc ttctatcgcc 5760
 ttcttgacga gttcttctga gcgggactct ggggttcgaa tcctaccagg gattggctga 5820
 ctgatgagtc cgtgaggac aaacgaaaaa catatggtac 5860

<210> 6

55 <211> 4610

EP 1 035 209 A1

<212> DNA

<213> Influenza virus

<400> 6

```

5  gaggcatttc agtcagttgc tcaaggtacc aaaatgaaca ctcaaatect ggttttcgcc 60
   cttgcggcag tcatccccac aaatgcagac aaaatttgtc ttggacatca tgctgtatca 120
   aatggcacca aagtaaacac actcactgag agaggagtag aagtrgtcaa tgcaacggaa 180
   acagtggagc ggacaaacat ccccaaaatt tgctcaaaag ggaaaagaac cactgatctc 240
   ggccaatgcg gactgttagg gaccattacc ggaccacctc aatgcgacca atttctagaa 300
10  ttttcagctg atctataaat cgagagacga gaaggaaatg atgtttgtta cccggggaag 360
   tttgttaatg aagaggcatt gcgacaaatc ctcagaggat cagggtgggat tgacaaaqaa 420
   acaatgggat tcacatatag tggaataagg accaacggaa caactagtgc atgtagaaga 480
   tcagggtcct cattctatgc agaaatggag tggctcctgt caaatacaga caatgcttct 540
   ttcccacaaa tgacaaaatc atacaaaaac acaggggagag aatcagctct gatagtctgg 600
15  ggaatccacc attcaggatc aaccaccgaa cagaccaaac tataatggag tggaataaaa 660
   ctgataacag tcgggagttc caaatatcat caatcttttg tgccgagtc aggaacacga 720
   ccgcagataa atggccggtc cggacggatt gattttcatt ggttgatctt ggatcccaat 780
   gatacagtta ctttttagttt caatggggct ttcatagtct caaatcgtgc cagcttcttg 840
   aggggaaagt ccatggggat ccagagcgat gtgcaggttg atgctaattg cgaaggggaa 900
20  tgctaccaca gtggagggac tataacaagc agattgcctt ttcaaaacat aaatagcaga 960
   gcagttggca aatgccccag atatgtaaaa caggaaaagt tattattggc aactgggatg 1020
   aagaacggtc ccgaaccttc caaaaaaagg aaaaaaagag gcctgtttgg tgctatagca 1080
   ggggtttattg aaaatggttg ggaaggctcg gtgcaggggt ggtaagggtt caggcatcag 1140
   aatgcacaag gagaagggaac tgcagcagac taaaaagca cccaatcggc aattgatcag 1200
25  ataaccggaa agttaaatag actcattaag aaaaccaacc agcaatttga gctaatagat 1260
   aatgaattca ctgaagtgga aaagcagatt ggcaatttaa ttaactggac caaagactcc 1320
   atcacagaag tatggtctta caatgctgaa cttcttgttg caatggaaaa ccagcacact 1380
   attgatttgg ctgattcaga gatgaacaag ctgtatgagc gagtgaggaa acaattaagg 1440
   gaaaatgctg aagaggatgg cactgggttc tttgaaattt ttcataaatg tgacgatgat 1500
30  tgtatggcta gtataaggaa caatacttat gatcacagca aatacagaga agaagcgatg 1560
   caaaatagaa tacaaattga cccagtcaaa ttgagtagtg gctacaaaga tgtgatactt 1620
   tgggttagct tcggggcatc atgctttttg cttcttgcca ttgcaatggg ccttgttttc 1680
   atatgtgtga agaacggaaa catgcggtgc actatttga tataggtttg gaaaaaaca 1740
   ccccttgttt ctactcccc ccaacttcgg aggtcgacca gtactccggg cgaaactttg 1800
35  tttttttttt ttccccgat gctggaggtc gaccagatgt ccgaaagtgt ccccccccc 1860
   ccccccccc ggcgcggaac ggcggggcca ctctggactc tttttttttt tttttttttt 1920
   ttttttgggg atcgccgct agcttctgtt ttggcggatg agagaagatt ttcagcctga 1980
   tacagattaa atcagaacgc agaagcggtc tgataaaaaca gaatttgcct ggcggcagta 2040
   gcgcggtggt cccacctgac cccatgccga actcagaagt gaaacgccgt agcgccgatg 2100
40  gtagtggtgg gtctcccat gcgagagtag ggaactgcc a ggcatcaaat aaaacgaaag 2160
   gctcagtcga aagactgggc ctttcgtttt atctgttgtt tgcgttgaa cgctctcctg 2220
   agtaggacaa atccgcccgg agcggatttg aacgttgcca agcaacggcc cggagggtgg 2280
   cgggcaggac gcccgccata aactgccagg catcaaatta agcagaaggc catcctgacg 2340
   gatggccttt ttgcgtttct aaaaactctt ttgtttattt ttctaaatc attcaaatat 2400
45  gtatccgctc atgagacaat aacctgata aatgcttcaa taatattgaa aaaggaagag 2460
   tatgagtatt caacatttcc gtgtcgccct tattcccttt tttgcggcat tttgccttcc 2520
   tgtttttgct caccagaaa cgctggtgaa aytaaaagat gctgaagatc agttgggtgc 2580
   acgagtgggt tacatcgaac tggatctcaa cagcggtaag atccttgaga gttttcgccc 2640
   cgaagaacgt tttccaatga tgagcacttt taaagtctg ctatgtggcg cggattattc 2700
50  ccgtgttgac gccgggcaag agcaactcgg tcgcgcgata cactattctc agaatgactt 2760
   ggttgagtac tcaccagtca cagaaaagca tcttacggat ggcatgacag taagagaatt 2820
   atgcagtgtc gccataacca tgagtataa cactgcggcc aacttacttc tgacaacgat 2880
   cggaggaccg aaggagctaa ccgctttttt gcacaacatg ggggatcatg taactcgcc 2940
   tgatcgttgg gaaccggagc tgaatgaagc cataccaaac gacgagcgtg acaccacgat 3000
55  gcctgtagca atggcaacaa cgttgcgcaa actattaact ggcgaactac ttactctagc 3060

```

5 ttccccgcaa caattaatag actggatgga ggccgataaa gttgcaggac cacttctgcg 3120
 ctccggccctt ccggctggct gggtttattgc tgataaatct ggagccgggtg agcgtgggtc 3180
 tcgcggtatc attgcagcac tggggccaga tggtaagccc tcccgatcgc tagttatcta 3240
 cacgacgggg agtcaggcaa ctatggatga acgaaataga cagatcgctg agataggtgc 3300
 ctcaactgatt aagcattggc aactgtcaga ccaagtttac tcatatatac tttagattga 3360
 tttaaaactt catttttaat ttaaaaggat ctaggatgaag atcctttttg ataattctat 3420
 gacaaaaatc ccttaacgtg agtttttcgtt ccactgagcg tcagaccccc tagaaaaagat 3480
 caaaggatct tcttgagatc ctttttttct gcgcgtaatc tgctgcttgc aaacaaaaaa 3540
 10 accaccgcta ccagcgggtg tttgtttgcc ggatcaagag ctaccaatc tttttccgaa 3600
 ggtaactggc ttcagcagag cgcagatacc aaatactgtc cttctagtgt agccgtagt 3660
 aggccaccac ttcaagaact ctgtagcacc gcctacatac ctgctctgc taatcctgtt 3720
 accagtggct gctgccagt gcgataagtc gtgtcttacc ggggtggact caagacgata 3780
 gttaccggat aaggcgcagc ggtcgggctg aacggggggg tctgtcacac agcccagctt 3840
 ggagcgaacg acctacaccg aactgagata cctacagcgt gagctatgag aaagcgcac 3900
 15 gcttccccgaa gggagaaaag cggacaggta tccggtaagc ggagggtcg gaacaggaga 3960
 gcgcacgagg gagcttccag ggggaaacgc ctggtatctt tatagtcctg tcgggtttcg 4020
 ccacctctga cttgagcgtc gatttttgtg atgtcgtca gggggcgga gcctatggaa 4080
 aaacgccagc aacgcggcct ttttacggtt cctggccttt tgctggcctt ttgtcacat 4140
 gttctttcct gcgttatccc ctgattcatt aatgcaggtc acgatccttt ctggcgagtc 4200
 20 cccgtgcgga gtcggagagc gctccctgag cgcgtgcggc ccgagaggtc gcgcctggcc 4260
 ggccttcggg cctcgtgtg tcccggtcgt agggggggcc ggccgaaaat gcttccggt 4320
 cccgctctgg agacacgggc cggcccccctg cgtgtggcac gggcgcccg gagggcgtcc 4380
 ccggcccggc yctgtcccg cgtgtgtcct ggggttgacc agaggggccc gggcgctccg 4440
 tgtgtggctg cgatgggtgg gtttttgggg acaggtgtcc gtgtccgtgt cgcgcgtcgc 4500
 25 ctgggcccggc ggctgtggctg gtgacgcgac ctcccgggcc cgggggagggt atatctttcg 4560
 ctccgagtcg gcattttggg ccgccggggt attagtagaa acaggggtac 4610

<210> 7

<211> 3558

30 <212> DNA

<213> Influenza virus

<400> 7

35 tattagtaga aacagggtat tttttattct agtacattac gccccgccct gccactcatc 60
 gcagtactgt tgtaattcat taagcattct gccgacatgg aagccatcac agacggcatg 120
 atgaacctga atcgccagcg gcatcagcac cttgtgcct tgcgataaat atttgcccat 180
 ggtgaaaacg gggggaaga agttgtccat attggccacg tttaaatcaa aactggtgaa 240
 actcaccag ggattggctg agacgaaaaa catattctca ataaaccctt tagggaaata 300
 ggccagggtt tcaccgtaac agccacatc ttgcgaatat atgtgtagaa actgccggaa 360
 40 atcgtcgtgg tattcactcc agagcgatga aaacggttca gtttgcctcat ggaaaacggg 420
 gtaacaaggg tgaacactat cccatatac cagctcaccg tctttcattg ccatacggaa 480
 ttccggatga gcattcatca ggccggcaag aatgtgaata aaggccggat aaaacttgtg 540
 cttatttttc tttacggtct ttaaaaaggc cgtaatatcc agctgaacgg tctggttata 600
 ggtacattga gcaactgact gaaatgcctc aaaatgttct ttacgatgcc attgggatat 660
 45 atcaacggtg gtatatccag tgattttttt ctccatgatt atggccatta ccctgtttc 720
 tactcccccc caacttcgga ggtcgaccag tactccgggc gaaactttgt tttttttttt 780
 tcccccgatg ctggaggctg accagatgtc cgaaagtgtc cccccccccc ccccccccg 840
 gcgcggaacg gcggggccac tctggactct tttttttttt tttttttttt tttttgggga 900
 tcggcgceta gcttctgttt tggcggatga gagaagattt tcagcctgat acagattaaa 960
 50 tcagaacgca gaagcggctc gataaaacag aatttgcctg gcggcagtag cgcggtggtc 1020
 ccactgacc ccacggcga ctcagaagtg aaacgcgta gcgccgatg tagtgtggg 1080
 tctcccatg cgagagtagg gaactgccag gcatcaaata aaacgaaagg ctcatcgaa 1140
 agactgggcc tttcgtttta tctgtgtgtt gtcggtgaac gctctcctga gtaggacaaa 1200
 tccgcgggga gcggttttga acgttgcgaa gcaacggccc ggagggtggc gggcaggaa 1260
 55 ccgccataa actgccaggc atcaaattaa gcagaaggcc atcctgacgg atggcctttt 1320

5 tgcgtttcta caaactcttt tgtttatttt tctaaataca ttcaaataatg tatccgctca 1380
 tgagacaata accctgataa atgcttcaat aatattgaaa aaggaagagt atgagtattc 1440
 aacatttccg tgcgcctt attccctttt ttgcggcatt ttgccttcct gt ttttlyc 1500
 acccagaaac gctggtgaaa gtaaaagatg ctgaagatca gttgggtgca cgagtgggtt 1560
 acatcgaact ggatctcaac agcggtaaga tccttgagag ttttcgcccc gaagaacgtt 1620
 ttccaatgat gagcactttt aaagtctcgc tatgtggcgc ggtattatcc cgtgttgacg 1680
 ccgggcaaga gcaactcggg cgcgcgcatc actattctca gaatgacttg gttgagtact 1740
 caccagtcac agaaaagcat cttacggatg gcatgacagt aagagaatta tgcagtgcct 1800
 10 ccataaccat gaggataaac actgcggcca acttacttct gacaacgac ggaggaccga 1860
 aggagctaac cgcttttttg cacaacatgg gggatcatgt aactcgctt gatcgttggg 1920
 aaccggagct gaataagcc ataccaaacg acgagcgtga caccacgatg cctgtagcaa 1980
 tggcaacaac gttgcgcaaa ctattaactg gcgaactact tactctagct tcccggcaac 2040
 aattaataga ctggatggag gcggataaag ttgcaggacc acttctgcgc tcggcccttc 2100
 cggtcgtgct gtttattgct gataaatctg gagccggtga gcgtgggtct cgcggtatca 2160
 15 ttgcagcact ggggccagat ggtaagccct cccgtatcgt agttatctac acgacgggga 2220
 gtcaggcaac tatggatgaa cgaaatagac agatcgctga gatagggtgcc tactgatta 2280
 agcattggta actgtcagac caagtttact catatatact ttagattgat ttaaaacttc 2340
 atttttaatt taaaaggatc taggtgaaga tcctttttga taatctcatg accaaaatcc 2400
 20 cttaacgtga gttttcgttc cactgagcgt cagaccccg agaaaagatc aaaggatctt 2460
 cttgagatcc tttttttctg cgcgtaatct gctgcttgca aacaaaaaaa ccaccgctac 2520
 cagcgggtgt ttgtttgccc gatcaagagc taccaactct ttttccgaag gtaactggct 2580
 tcagcagagc gcagatacca aatactgtcc ttctagtga gccgtagtta ggccaccact 2640
 tcaagaactc tgtagcacc cctacatacc tcgctctgct aatcctgta ccagtggctg 2700
 ctgccagtgg cgataagtcg tgccttaccg ggttggactc aagacgatag ttaccggata 2760
 25 aggcgcagcg gtcgggctga acgggggggt cgtgcacaca gccagcttg gacgcaacga 2820
 cctacaccga actgagatac ctacagcgtg agctatgaga aagcgccacg cttcccgaag 2880
 ggagaaaggc ggacaggat ccggttaagcg gcagggtcgg aacaggagag cgcacgaggg 2940
 agcttccagg gggaaacgcc tggatatctt atagtcctgt cgggtttcgc cacctctgac 3000
 ttgagcgtcg atttttgtga tgcctgcag gggggcggag cctatggaaa aacgccagca 3060
 30 acgcggcctt tttacggttc ctggcctttt gctggccttt tgctcacatg ttctttcctg 3120
 cgttatcccc tgattcatta atgcaggatc cgtcccttc tggcgagtc ccgtgcggag 3180
 tcggagagcg ctccctgagc gcgtgcggcc cgagaggctc cgcctggccg gccttcggtc 3240
 cctcgtgtgt cccggtcgta ggagggggcg gccgaaaatg cttccgctc ccgctctgga 3300
 gacacggggc ggccccctgc gtgtggcac ggcggccggg agggcgctcc cggccccggc 3360
 35 ctgctcccgc gtgtgtcctg gggttgacca gagggccccg ggcgctccgt gtgtggctgc 3420
 gatggtggcg tttttgggga cagggtgcgc tgcctgtgc gcgcgtcgcc tgggccccg 3480
 gcgtggtcgg tgacgcgacc tcccggcccc gggggaggta tatctttcgc tccgagtcgg 3540
 cattttgggc cgccgggt 3558

40 <210> 8
 <211> 4343
 <212> DNA
 <213> Influenza virus

45 <400> 8
 ctttctggcg agtccccgtg cggagtcgga gacgcctccc tgagcgcgtg cggccccgaga 60
 ggtcgcgcct ggccggcctt cggctccctcg tgtgtcccgg tcgtaggagg ggccggccga 120
 aaatgcttcc ggctcccgtc ctggagacac gggccggccc cctgcgtgtg gcacgggcgg 180
 ccgggagggc gtccccggcc cggcgctgct cccgcgtgtg tctgggggtt gaccagaggg 240
 50 ccccggggcg tccgtgtgtg gctgcgatgg ttggcgtttt ggggacaggt gtccgtgtcc 300
 gtgtcgcgcg tcgcctgggc cggcggcgtg gtcggtgac cgacctccc gccccggggg 360
 aggtatatct ttgcgtccga gtcggcattt tgggccccg ggttattagt agaaacaggg 420
 ttttttttat actagtaagc tcgaaggagt ccaccatgag taaaggagaa gaacttttca 480
 ctggagttgt cccaattctt gttgaattag atgggtgatg taatgggcac aaattttctg 540
 55 tcagtggaga ggggtgaagg gatgcaacat acggaaaact tacccttaaa tttatttcga 600

ctactggaaa actacctgtt ccatggccaa cacttgctac tactttcact tatgggtgttc 660
 aatgcttttc aagataccca gatcatatga aacagcatga ctttttcaag agtgccatgc 720
 ccgaagggtta tgtacaggaa agaactatat ttttcaaaga tgacgggaac tacaagacac 780
 gtgctgaagt caagtttgaa ggtgatcccc ttgttaatag aatcgagtta aaagggtattg 840
 atttttaaaga agatggaaac attcttggtac acaaattgga atacaactat aactcacaca 900
 atgtatacat catggctgac aagcagaaga acggaatcaa ggccaacttc aagaccgcc 960
 acaacatcga ggacggcggc gtgcagctgg ccgaccacta ccagcagaac accccaattg 1020
 gcgatggccc tgctctttta ccagacaacc attacctgtc cacacaatct gccctttcga 1080
 aagatcccaa cgaaaagaga gaccacatgg tcttcttga gtttgtaaca gctgctggga 1140
 ttacacatgg catggatgaa ctatacaagg gatcccatca ccatcaccat cactaagctc 1200
 catggctctag atatctagta cattacgccc cgccctgcca ctcatcgag tactgttga 1260
 attcattaag cattctgccc acatggaagc catcacagac ggcatgatga acctgaatcg 1320
 ccagcggcat cagcaccttg tcgcttgcg tataatattt gcccatggtg aaaacggggg 1380
 cgaagaagtt gtccatattg gccacgttta aatcaaaact ggtgaaactc acccagggat 1440
 tggcactcac aaagaacatg ttctcgatga atcctttagg gaagtaggcc aggttttcac 1500
 cgtaacacgc cacatcttgc gaatatatgt gtagaaactg ccggaaatcg tcgtggtatt 1560
 cactccagag cgatgaaaac gtttcagttt gctcatggaa aacggtgtaa caagggtgaa 1620
 cactatccca tatcaccagc tcaccgctct tcatggccat acggaattcc ggatgagcat 1680
 tcatcaggcg ggcaagaatg tgaataaagg ccggataaaa cttgtgctta tttttcttta 1740
 cggctcttaa aaaggccgta atatccagct gaacggctcg gttataggta cattgagcaa 1800
 ctgactgaaa tgctcaaaa tgttctttac gatgccattg ggatatatca acggtggtat 1860
 atccagtgat tttttctcc atgattatgc aaaaaatacc cttgtttcta ctcccccca 1920
 acttcggagg ctccagcagta ctccggcgga aactttgttt tttttttttt ccccgatgct 1980
 ggaggctgac cagatgtccc aaagtgtccc ccccccccc cccccccgc gcggaacgc 2040
 gggggccactc tggactcttt tttttttttt tttttttttt tttggggatc ggccgctagc 2100
 ttctgttttg gcggatgaga gaagattttc agcctgatac agattaaatc agaacgcaga 2160
 agcggctcga taaaacagaa ttgctctggc ggcagtagcg cgggtggtccc acctgacccc 2220
 atgccgaact cagaagtga acgcccgtagc gccgatggta gtgtggggtc tccccatgcg 2280
 agagtaggga actgccaggc atcaaaataaa acgaaaggct cagtcgaaag actgggcctt 2340
 tcgttttatc tgttgtttgt cgggtgaacgc tctcctgagt aggacaaatc cgccgggagc 2400
 ggatttgaac gttgcgaagc aacggcccg aggggtggcg gcaggacgcc cgccataaac 2460
 tgccaggcat caaatgaagc agaaggccat cctgacggat ggcctttttg cgtttctaca 2520
 aactcctttg ttttttttc taaatacatt caaatatgta tccgctcatg agacaataac 2580
 cctgataaat gcttcaataa tattgaaaaa ggaagagtat gagtattcaa catttccgtg 2640
 tcgccccttat tccctttttt gcggcatttt gccttctgt ttttgctcac ccagaaacgc 2700
 tgggtgaaagt aaaagatgct gaagatcagt tgggtgcacg agtgggttac atcgaactgg 2760
 atctcaacag cggtaagatc cttgagagtt ttgcctccga agaacgtttt ccaatgatga 2820
 gcacttttaa agttctgcta tgtggcgcg tattatcccg tgttgacgcc gggcaagagc 2880
 aactcggctg ccgcatacac tattctcaga atgacttggg tgagtactca ccagtcacag 2940
 aaaagcatct tacggaaggc atgacagtaa gagaattatg cagtgtctgc ataaccatga 3000
 gtgataacac tgcggccaac ttacttctga caacgatcgg aggaccgaag gagctaaccg 3060
 cttttttgca caacatgggg gatcatgtaa ctgccttga tcgttgggaa ccggagctga 3120
 atgaagccat accaaacgac gacggtgaca ccacgatgcc tgtagcaatg gcaacaacgt 3180
 tgcgcaaat attaaatggc gaactactta ctctagcttc ccggcaacaa ttaatagact 3240
 ggatggaggc ggataaagtt gcaggaccac ttctgcgctc ggcccttccg gctggctggt 3300
 ttattgctga taaatctgga gccggtgagc gtgggtctcg cggtatcatt gcagcactgg 3360
 ggccagatgg taagccctcc cgtatcgtag ttatctacac gacggggagt caggcaacta 3420
 tggatgaacg aaatagacag atcgctgaga taggtgcctc actgattaag catttgtaac 3480
 tgtcagacca agtttactca tatatacttt agattgattt aaaacttcat ttttaattta 3540
 aaaggatcta ggtgaagatc ctttttgata atctcatgac caaaatccct taacgtgagt 3600
 ttctggtcca ctgagcglca gacccctag aaaagatcaa aggatcttcl lgatctctt 3660
 tttttctgcg cgtaatctgc tgcttgcaaa caaaaaaac accgctacca gcgggtggtt 3720
 gtttgccgga tcaagagcta ccaactcttt ttccgaaggt aactggcttc agcagagcgc 3780
 agataccaaa tactgtctct ctagtgtagc cgtagttagg ccaccacttc aagaactctg 3840
 tagcaccgcc tacatacctc gctctgctaa tctgttacc agtggctgct gccagtggcg 3900

EP 1 035 209 A1

ataagtcgtg tcttaccggg ttggactcaa gacgatagtt accggataag gcgcagcggg 3960
 cgggctgaac ggggggttcg tgcacacagc ccagcttggg gcgaacgacc tacaccgaac 4020
 tgagatacct acagcgtgag ctatgagaaa gcgccacgct tcccgaaggg agaaagggcg 4080
 5 acaggtatcc ggtaagcggc agggctcgga caggagagcg cagcagggag ctccagggg 4140
 gaaacgcctg gtatctttat agtctgtcg ggtttcgcca cctctgactt gagcgtcgat 4200
 ttttgtgatg ctcgtcaggg gggcggagcc tatggaaaaa cgccagcaac gcggcctttt 4260
 tacggttcct ggccttttgc tggccttttg ctacatgtt ctttctcgcg ttatccctg 4320
 attcattaat gcaggtcacg atc . 4343

10 <210> 9
 <211> 3888
 <212> DNA
 <213> Influenza virus

15 <400> 9
 cccaaaaaaa aaaaaaaaaa aaaaaaaaaa agtccagagt ggccccgccc ttccgcgccc 60
 gggggggggg ggggggggga cactttcggg catctggtcg acctccagca tcgggggaaa 120
 aaaaaaaaaa aaagtttcgc ccggagtact ggtcgacctc cgaagttagg ggggagtaga 180
 20 aacagggtag ataactcact actgagtgc atccacatcg cgagcgcgcg taatacgact 240
 cactataggg cgaattgggt accgggcccc cctcgaggt cgacggtatc gataagcttc 300
 gacgagatct tcaggagcta aggaagctaa aatggagaaa aaaatcactg gatataccac 360
 cgttgatata tcccaatggc atcgtaaaga acattttgag gcatttcagt cagttgctca 420
 atgtacctat aaccagaccg ttcagctgga tattacggcc tttttaaaga ccgtaaagaa 480
 aaataagcac aagtttttat cggcctttat tcacattctt gcccgcctga tgaatgctca 540
 25 tccggaattc cgtatggcaa tgaaagacgg tgagctggtg atatgggata gtgttcaccc 600
 ttgttacacc gttttccatg agcaaaactga aacgttttca tcgctctgga gtgaatacca 660
 cgacgatttc cggcagtttc tacacatata ttcgcaagat gtggcgtgtt acggtgaaaa 720
 cctggcctat ttccctaaag ggtttattga gaatatgtt ttcgtctcag ccaatccctg 780
 ggtgagtttc accagttttg atttaaacgt ggccaatatg gacaacttct tcgccccctg 840
 30 tttcaccatg ggcaaatatt atacgcaagg cgacaagggt ctgatgccgc tggcgattca 900
 ggttcatcat gccgtttgtg atggcttcca tgtcggcaga atgcttaatg aattacaaca 960
 gtactgcgat gagtggcagg gcggggcgta atttttttaa ggcagttatt ggtgccctta 1020
 aacgcctggt gctacgcctg aataagtgat aataagcgga tgaatggcag aaattcgtcg 1080
 aagcttgata tcgaattcct gcagcccggg ggatccacta gttctagagc ggccgcccac 1140
 35 gcggtggagc tccagctttt gttcccttta gtgagggtta attgcgcgca ggccctagcta 1200
 ggtaaagaaa aatacccttg attcttctaa taaccggcg gcccaaatg ccgactcgga 1260
 gcgaaagata tacctcccc ggggcccggga ggtcgcgtca ccgaccagc cgccggccca 1320
 ggcgacgcgc gacacggaca cctgtcccca aaaacgccac catcgcagcc acacacggag 1380
 cgccccgggc cctctggtca accccaggac acacgcggga gcagcgcggg gccggggacg 1440
 40 ccctcccggc cgcccggtgc acacgcaggg gccgggcccg tgtctccaga gcgggagccg 1500
 gaagcatttt cggccggccc ctctacgac cgggacacac gagggaccga aggccggcca 1560
 ggcgcgacct ctggggcgc acgcgcgctc agggagcgct ctccgactcc gcacggggac 1620
 tcgccagaaa ggatcgtgac ctgcattaat gaatacgggg ataacgcagg aaagaacatg 1680
 tgagcaaaaag gccagcaaaa ggccaggaac cgtaaaaagg ccgcgttgct ggcgttttc 1740
 45 cataggctcc gccccctga cgagcatcac aaaaatcgac gctcaagtca gaggtggcga 1800
 aaccggacag gactataaag ataccaggcg tttccccctg gaagctccct cgtgcgctct 1860
 cctgttccga ccctgccgct taccggatac ctgtccgctt ttctcccttc gggaagcgtg 1920
 gcgctttctc atagctcacg ctgtaggtat ctgagttcgg ttaggtcgt tcgctccaag 1980
 ctgggtctgt tgcacgaacc ccccgttcag cccgaccgct gcgccttctc cggttaactat 2040
 50 cgtcttgagt ccaaccgggt aagacacgac ttatcgccac tggcagcagc cactggtaac 2100
 aggattagca gagcagggta tgtaggcggg gctacagagt tcttgaagt gtggcctaac 2160
 tacggctaca ctagaaggac agtatattgt atctgcgctc tgctgaagcc agttaccttc 2220
 ggaaaaagag ttggtagctc ttgatccggc aaacaaacca ccgctggtag cgggtggtttt 2280
 tttgtttgca agcagcagat tacgcgcaga aaaaaggat ctcaagaaga tcctttgatc 2340
 55 ttttctacgg ggtctgacgc tcagtggaaac gaaaactcac gtaagggat tttggtcatg 2400

EP 1 035 209 A1

agattatcaa aaaggatctt cacctagatc cttttaaatt aaaaatgaag ttttaaatca 2460
 atctaaagta tatatgagta aacttggctt gacagttacc aatgcttaat cagtgaggca 2520
 cctatctcag cgatctgtct atttcgttca tccatagtgt cctgactccc cgtcgtgtag 2580
 5 ataactacga tacgggaggg cttaccatct ggccccagtg ctgcaatgat accgcgagac 2640
 ccacgctcac cggctccaga tttatcagca ataaaccagc cagccggaag ggccgagcgc 2700
 agaagtggc ctgcaacttt atccgcctcc atccagtcta ttaattgttg ccgggaagct 2760
 agaytaagta gttcgcagc taatagtgtg cgcaacgttg ttgccattgc tacaggcacc 2820
 gtgggtgtcac gtcgtcgtt tggatggct tcattcagct ccggttccca acgatcaagg 2880
 10 cgagttacat gatccccat gttgtgcaaa aaagcgggta gtccttcgg tcttcggatc 2940
 gttgtcagaa gtaagtggc cgcagtgtta tcactcatgg ttatggcagc actgcataat 3000
 tctcttactg tcatgccatc cgtaagatgc tttctgtga ctggtgagta ctcaaccaag 3060
 tcattctgag aatagtgtat gcggcgaccg agttgctctt gcccgcgctc aacacgggat 3120
 aataccgcgc cacatagcag aactttaaaa gtgctcatca ttggaaaacg tcttcggggg 3180
 cgaaaactct caaggatctt accgctgttg agatccagtt cgatgtaacc cactcgtgca 3240
 15 cccaactgat cttcagcatc ttttactttc accagcgttt ctgggtgagc aaaaacagga 3300
 aggcaaaatg ccgcaaaaaa gggaataagg gcgacacgga aatgttgaat actcatactc 3360
 ttcctttttc aatattattg aagcatttat cagggttatt gtctcatgag cggatacata 3420
 tttgaatgta tttagaaaaa taaacaaaag agtttgtaga aacgcaaaaa ggccatccgt 3480
 20 caggatggcc ttctgcttaa ttgatgctt ggcagtttat ggccggcgctc ctgcccgcca 3540
 ccttcggggc cgttgccttc caacgttcaa atccgctccc ggccgatttg tcttactcag 3600
 gagagcgttc accgacaaac aacagataaa acgaaaggcc cagtctttcg actgagcctt 3660
 tcgtttttatt tgatgccttg cagttcccta ctctcgcatg gggagacccc aactaccat 3720
 cggcgctacg gcgtttcact tctgagttcg gcatggggtc aggtgggacc accgcgctac 3780
 25 tgccgccagg caaattctgt tttatcagac cgcttctcg tttctgattta atctgtatca 3840
 ggctgaaaat cttctctcat ccgcaaaaac agaagctagc ggccgac 3888

30

35

40

45

50

55

Claims

1. A recombinant influenza virus for high-yield expression of incorporated foreign gene(s), which is genetically stable in the absence of any helper virus; wherein
 - (a) at least one of the regular viral RNA segments has been exchanged for a vRNA encoding a foreign gene; and/or
 - (b) at least one of the regular viral RNA segments is an ambisense RNA molecule, containing one of the standard viral genes in sense orientation and a foreign, recombinant gene in anti-sense orientation, or *vice versa*, in overall convergent arrangement.
2. The recombinant virus according to claim 1, wherein in the ambisense RNA molecule said foreign recombinant gene is covalently bound to one of the viral genes, while the original vRNA segment coding for the same gene is deleted from the recombinant virus by way of specific ribozyme cleavage.
3. The recombinant virus according to claims 1 and 2 in which one or both of the standard glycoproteins hemagglutinin and neuraminidase have been exchanged into foreign glycoprotein(s) or into fusion glycoproteins consisting of an anchor segment derived from hemagglutinin and an ectodomain obtained from the foreign source, viral or cellular, or in which such recombinant glycoprotein has been inserted as a third molecular species in addition to the remaining standard components.
4. The recombinant virus according to claims 1 to 3, in which the terminal viral RNA sequences, which are active as the promoter signal, have been varied by nucleotide substitutions in up to five positions, resulting in improved transcription rates of both the vRNA promoter as well as the cRNA promoter as present in the complementary sequence.
5. The recombinant virus according to claims 1 to 4, in which the foreign gene(s) in ambisense covalent junction with viral gene(s) code for proteins and/or glycoproteins which are secreted from cells infected with the recombinant virus.
6. The recombinant virus according to claims 1 to 4, in which the foreign gene(s) in ambisense covalent junction with viral gene(s) code for proteins or artificial polypeptides designed to support an efficient presentation of inherent epitopes at the surface of (abortively) infected cells, for stimulation of a B cell and/or T cell response.
7. A method for the production of recombinant influenza viruses as defined in claims 1 to 6 comprising
 - (a) RNA polymerase I synthesis of recombinant vRNAs *in vivo*, in antisense, sense or ambisense design,
 - (b) followed by infection with an influenza carrier strain constructed to include flanking ribozyme target sequences in at least one of its viral RNA segments, and
 - (c) thereafter selective vRNA inactivation through ribozyme cleavage.
8. A method of constructing influenza carrier strains carrying one or more ribozyme target sites (of type one) in vRNA flanking positions comprising
 - (a) RNA polymerase I synthesis of recombinant vRNAs *in vivo*, carrying two different 3' promoter sequences in tandem, which are separated by a second type of ribozyme target sequence, and which carry the said internal ribozyme target sites of type one;
 - (b) followed by infection of an influenza wildtype strain;
 - (c) thereafter amplification through simple steps of viral propagation; and
 - (d) finally isolation through removal of their external 3' promoter sequence by ribozyme cleavage through infection of cells expressing ribozyme type 2, followed by plaque purification.
9. A ribozyme-sensitive influenza carrier strain obtainable by the method of claim 8.
10. A pharmaceutical composition comprising a recombinant virus according to claims 1 to 6.
11. Use of a recombinant virus according to claims 1 to 6 for preparing a medicament for vaccination purposes.

12. The use according to claim 11, wherein the medicament

- (a) is suitable against influenza and/or against other infections;
- (b) is present in form of inactivated preparations; and/or
- (c) is present in form of live recombinant (attenuated) viruses.

13. Use of a recombinant virus according to claims 1 to 6 as vector systems in somatic gene therapy, for transfer and expression of foreign genes into cells (abortively) infected by such viruses, either in *ex vivo* or *in vivo* application schemes.

14. A method for the production of proteins or glycoproteins which comprises utilizing a recombinant influenza virus according to claims 1 to 6 as expression vector.

15. The method of claim 14, wherein the production is performed in cell culture cells or in fertilized chicken eggs.

FIG. 1




FIG. 2





FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9

cRNA-promoter





FIG. 11




FIG. 12




FIG. 13



FIG. 14




FIG. 15




FIG. 16




FIG. 17




FIG. 18




FIG. 19





FIG. 20





European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention shall be considered, for the purposes of subsequent proceedings, as the European search report

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION
X	DE 197 09 512 A (HOBOM GERD PROF DR DR) 10 September 1998 (1998-09-10)	1,3,4, 7-14	C12N15/86 C12N7/01
Y	* the whole document *	2,5,6,15	C12N5/10 A61K39/00 A61K39/145 A61K48/00
X	WO 91 03552 A (SINAI SCHOOL MEDICINE) 21 March 1991 (1991-03-21)	1,10-15	
Y	* figure 11; example 7 *	2,5,6	
Y	TAKASE H. ET AL.: "Antibody responses and protection in mice immunized orally against influenza virus." VACCINE, vol. 14, no. 17/18, 1996, pages 1651-1656, XP002110225 * page 1652, left-hand column, paragraph 1 *	15	
X	ZHOU Y. ET AL.: "Membrane-anchored incorporation of a foreign protein in recombinant Influenza virions." VIROLOGY, vol. 246, 20 June 1998 (1998-06-20), pages 83-94, XP002110226 * the whole document *	1,10-14	
			TECHNICAL FIELDS SEARCHED
			C12N A61K
<p align="center">-/--</p>			
INCOMPLETE SEARCH			
<p>The Search Division considers that the present application, or one or more of its claims, does/do not comply with the EPC to such an extent that a meaningful search into the state of the art cannot be carried out, or can only be carried out partially, for these claims</p> <p>Claims searched completely :</p> <p>Claims searched incompletely :</p> <p>Claims not searched :</p> <p>Reason for the limitation of the search:</p> <p align="center">see sheet C</p>			
Place of search		Date of completion of the search	Examiner
THE HAGUE		26 July 1999	Mandl, B
CATEGORY OF CITED DOCUMENTS			
<p>X : particularly relevant if taken alone</p> <p>Y : particularly relevant if combined with another document of the same category</p> <p>A : technological background</p> <p>O : non-written disclosure</p> <p>P : intermediate document</p> <p>T : theory or principle underlying the invention</p> <p>E : earlier patent document, but published on, or after the filing date</p> <p>D : document cited in the application</p> <p>L : document cited for other reasons</p> <p>& : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03/82 (POIC07)



European Patent
Office

INCOMPLETE SEARCH
SHEET C

Application Number

EP 99 10 4519

Although claim 13 and claim 14, as far as an in vivo application is concerned, are directed to a method of treatment of the human/animal body (Article 52(4) EPC), the search has been carried out and based on the alleged effects of the compound/composition.

PARTIAL EUROPEAN SEARCH REPORT

Application Number
EP 99 10 4519

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
X	<p>PALESE P. ET AL.: "Negative-strand RNA viruses: Genetic engineering and applications." PROC. NATL. ACAD. SCI. U.S.A., vol. 93, October 1996 (1996-10), pages 11354-11358, XP000196755 * page 11354, right-hand column, last paragraph - page 11356, right-hand column, paragraph F *</p> <p style="text-align: center;">---</p>	1,10-14	
X	<p>ZOBEL A. ET AL.: "RNA polymerase I catalysed transcription of insert viral cDNA." NUCLEIC ACIDS RESEARCH, vol. 21, no. 16, 1993, pages 3607-3614, XP002110227 * page 3607, right-hand column, paragraph 2 * * page 3612, right-hand column, paragraph 2 - page 3613, left-hand column, line 1 * * page 3614, left-hand column, paragraph 2 *</p> <p style="text-align: center;">-----</p>	1,10	<p>TECHNICAL FIELDS SEARCHED</p>

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 99 10 4519

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

26-07-1999

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 19709512 A	10-09-1998	NONE	
WO 9103552 A	21-03-1991	US 5166057 A	24-11-1992
		AT 126272 T	15-08-1995
		AU 636916 B	13-05-1993
		AU 6411890 A	08-04-1991
		CA 2065245 A	01-03-1991
		DE 69021575 D	14-09-1995
		DE 69021575 T	14-12-1995
		DK 490972 T	30-10-1995
		EP 0490972 A	24-06-1992
		ES 2075901 T	16-10-1995
		GR 90100639 A	30-12-1991
		JP 5500607 T	12-02-1993
		PT 95124 A	18-04-1991
		US 5252289 A	12-10-1993
		US 5578473 A	26-11-1996
		US 5854037 A	29-12-1998
		US 5840520 A	24-11-1998
		US 5786199 A	28-07-1998
		US 5820871 A	13-10-1992

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82